



ELSEVIER



Available online at www.sciencedirect.com

ScienceDirect

indagationes
mathematicae

Indagationes Mathematicae 34 (2023) 259–273

www.elsevier.com/locate/indag

Special issue on the occasion of Jaap Korevaar’s 100-th birthday

A Fuglede type theorem for Fourier multiplier operators

Ben de Pagter^{a,*}, Werner J. Ricker^b

^a *Delft Institute of Applied Mathematics, Faculty EEMCS, Delft University of Technology, P.O. Box 5031, 2600 GA Delft, The Netherlands*

^b *Math.-Geogr. Fakultät, Katholische Universität Eichstätt-Ingolstadt, D-85072 Eichstätt, Germany*

Dedicated in admiration to Jaap Korevaar on the occasion of his 100th birthday

Abstract

Let E be a translation invariant Banach function space over an infinite compact abelian group G and M_φ be a Fourier multiplier operator (with symbol φ) acting on E . It is assumed that E has order continuous norm and that E is reflection invariant (which ensures that $\bar{\varphi}$ is also a multiplier symbol for E). The following Fuglede type theorem is established. Whenever T is a bounded linear operator on E satisfying $M_\varphi T = T M_\varphi$, then also $M_{\bar{\varphi}} T = T M_{\bar{\varphi}}$.

© 2022 The Author(s). Published by Elsevier B.V. on behalf of Royal Dutch Mathematical Society (KWG). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords: Compact abelian group; Translation invariant Banach function space; Fourier multiplier operator; Reflection invariance

1. Introduction

A classical result of B. Fuglede states if A is a bounded normal operator on a Hilbert space H and T is a bounded linear operator on H satisfying $TA = AT$, then $TA^* = A^*T$, [8]. More generally, C.R. Putnam showed if A and B are bounded normal operators on H such that $AT = TB$, then $A^*T = TB^*$, [16]. This more general version actually follows from the original Fuglede theorem, [3]. Various different proofs of Fuglede’s theorem are known; see, for example, [7], p. 934, [9,17,18].

An abstraction of the Fuglede theorem could be as follows. Let X be a Banach space and A belong to a subalgebra of the bounded linear operators on X which has an involution \sharp . Does it follow that $TA^\sharp = A^\sharp T$ whenever T is a bounded linear operator on X satisfying $TA = AT$? For $X = L^p(\mathbb{T})$, with $1 \leq p < \infty$ and \mathbb{T} the circle group, let $\mathcal{M}_p(\mathbb{T})$ denote the algebra

* Corresponding author.

E-mail addresses: b.depagter@tudelft.nl (B.d. Pagter), werner.ricker@ku.de (W.J. Ricker).

<https://doi.org/10.1016/j.indag.2022.10.005>

0019-3577/© 2022 The Author(s). Published by Elsevier B.V. on behalf of Royal Dutch Mathematical Society (KWG). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

of all bounded functions $\varphi : \mathbb{Z} \rightarrow \mathbb{C}$ having the property that, for every $f \in L^p(\mathbb{T})$ there exists $g \in L^p(\mathbb{T})$, necessarily unique, such that its Fourier transform satisfies $\hat{g} = \varphi \hat{f}$ on \mathbb{Z} . Denoting g by $M_\varphi^{(p)} f$, the bounded linear operator $M_\varphi^{(p)}$ so generated on $L^p(\mathbb{T})$ is called the p -multiplier operator corresponding to φ . It is well known that the complex conjugate $\bar{\varphi}$ of φ belongs to $\mathcal{M}_p(\mathbb{T})$ whenever $\varphi \in \mathcal{M}_p(\mathbb{T})$. Clearly $M_\varphi^{(p)} \mapsto M_{\bar{\varphi}}^{(p)}$ is an involution on the algebra of all p -multiplier operators on $L^p(\mathbb{T})$. In this setting the “Fuglede theorem” is indeed valid, [14].

The aim of this note is to give a far reaching generalization of the above result beyond the setting of L^p -spaces. The group \mathbb{T} can be replaced with any infinite compact abelian group G and the L^p -spaces can be replaced by the significantly larger class of *translation invariant Banach function spaces* E over G . However, for this class of spaces E three phenomena arise which do not occur for L^p -spaces over G . First, the space E need *not* contain $L^\infty(G)$. Second, the reflection of a function in E need not belong to E (which has the effect that $\varphi \in \mathcal{M}_E(G)$ need not imply that $\bar{\varphi}$ belongs to $\mathcal{M}_E(G)$) and third, the space of all continuous functions $C(G)$ on G , even if $C(G) \subseteq E$, need not be dense in E . However, for the very large and important class of spaces E which *do* have all three of these properties it is established in [Theorem 5.2](#) that the “Fuglede theorem” does hold.

Since the methods used rely heavily of the fact that the spaces E are all *Banach function spaces* over G (in particular, Banach lattices) and that certain non-trivial properties of E arising from translation invariance have to be developed and established, we need to include some sections to address these features. An attempt has been made to keep the note as self-contained as possible.

2. Preliminaries

In this section we collect together some definitions and facts concerning Banach function spaces. Let (Ω, Σ, μ) be a measure space. Since we are interested in the case where $\Omega = G$ is an infinite compact abelian group with normalized Haar measure μ , we will assume that $\mu(\Omega) = 1$ and that μ is atomless. We denote by $L^0(\mu)$ the space of all \mathbb{C} -valued, Σ -measurable functions (with the usual identification of functions which are equal μ -a.e. on Ω). Then $L^0(\mu)$ is a Dedekind complete complex Riesz space (or, vector lattice). Recall that a subset $A \subseteq L^0(\mu)$ is called an (order) *ideal* whenever A is a linear subspace of $L^0(\mu)$ with the property that $|f| \leq |g|$, with $f \in L^0(\mu)$ and $g \in A$, implies that $f \in A$.

Definition 2.1. A *Banach function space* (briefly B.f.s.) over (Ω, Σ, μ) is a Banach space $(E, \|\cdot\|_E)$, where $E \subseteq L^0(\mu)$ is an ideal and $\|\cdot\|_E$ is an absolutely monotone norm on E (that is, $\|f\|_E \leq \|g\|_E$ whenever $|f| \leq |g|$ in E).

Evidently, B.f.s.’ are complex (Dedekind complete) Banach lattices. Therefore, we can freely apply the general theory of Banach lattices to B.f.s.’ For the theory of Banach lattices we refer the reader to [1,13,21,23], and for the theory of B.f.s.’ to [22] (Chapter 15), [2]. The class of B.f.s.’ includes the L^p -spaces ($1 \leq p \leq \infty$), Orlicz spaces, Lorentz spaces, Marcinkiewicz spaces and many more. If E and F are two B.f.s.’ over (Ω, Σ, μ) satisfying $E \subseteq F$, then the natural embedding of E into F is continuous. Indeed, the embedding operator is linear and positive, and a positive linear operator on a Banach lattice is automatically continuous (see e.g. Theorem 83.12 in [23]).

Remark 2.2. In Chapter 15 of [22] and in the series of papers [12], B.f.s.’ are introduced via function norms. We briefly recall this approach. Denote by $M(\mu)$ the set of all (equivalence classes of) extended \mathbb{C} -valued measurable functions on Ω (that is, every $f \in M(\mu)$ is of the form $f = g + ih$ with $g, h : \Omega \rightarrow [-\infty, \infty]$ measurable). The set of all $[0, \infty]$ -valued functions in $M(\mu)$ is denoted by $M(\mu)^+$. A *function norm* ρ is defined to be a map $\rho : M(\mu)^+ \rightarrow [0, \infty]$ satisfying:

- (i) If $u \in M(\mu)^+$ and $\rho(u) = 0$, then $u = 0$;
- (ii) $\rho(\lambda u) = \lambda \rho(u)$ for all $\lambda \in \mathbb{R}^+$ and $u \in M(\mu)^+$;
- (iii) $\rho(u + v) \leq \rho(u) + \rho(v)$ for all $u, v \in M(\mu)^+$;
- (iv) $\rho(u) \leq \rho(v)$ whenever $u \leq v$ in $M(\mu)^+$.

If $u \in M(\mu)^+$ and $\rho(u) < \infty$, then $u(x) < \infty$ for μ -a.e. $x \in \Omega$ (i.e., $u \in L^0(\mu)^+$); see [22], Section 63, Theorem 1. Define

$$E_\rho = \{f \in M(\mu) : \rho(|f|) < \infty\}. \tag{1}$$

Then E_ρ is an ideal in $L^0(\mu)$. Setting $\|f\|_{E_\rho} = \rho(|f|)$ for all $f \in E_\rho$, the space $(E_\rho, \|\cdot\|_{E_\rho})$ is a normed space. If E_ρ is complete, then it is a B.f.s.

Conversely, if $(E, \|\cdot\|_E)$ is a B.f.s. and we set

$$\rho(u) = \begin{cases} \|u\|_E & \text{if } u \in E^+ \\ \infty & \text{if } u \notin E^+ \end{cases}, \quad u \in M(\mu)^+,$$

then ρ is a function norm and the space E is recovered via (1). This shows that the approach in [22] is equivalent to ours and so, all results from [22] may be used in our setting (see also [23], Section 112).

Given a B.f.s. $(E, \|\cdot\|_E)$ over (Ω, Σ, μ) , there exists a set $A_0 \in \Sigma$ with the property that $f = 0$ μ -a.e. on $\Omega \setminus A_0$ for each $f \in E$ and, for every $A \in \Sigma$ satisfying $A \subseteq A_0$ and $\mu(A) > 0$, there exists $B \in \Sigma$ such that $B \subseteq A$, $\mu(B) > 0$ and $\chi_B \in E$ (see [22], Section 72 or [23], Section 86). The set A_0 is uniquely determined (modulo μ -null sets) and is called the *carrier* of E . Moreover, there exists a sequence $(A_n)_{n=1}^\infty$ in Σ such that $A_n \subseteq A_0$ and $\chi_{A_n} \in E$ for all n , and $A_n \uparrow A_0$. Any B.f.s. that we encounter in this note has its carrier equal to Ω . In general, if $E \neq \{0\}$, then one can replace, without loss of generality, Ω by A_0 .

Recall that a B.f.s. $(E, \|\cdot\|_E)$ over (Ω, Σ, μ) has *order continuous norm* (briefly o.c.-norm) if $\|f_\tau\|_E \downarrow_\tau 0$ whenever $f_\tau \downarrow_\tau 0$ in E (that is, (f_τ) is a downwards directed net in the positive cone E^+ with infimum 0). Since E is Dedekind complete, it follows from Theorem 103.6 in [23] that E has o.c.-norm if and only if E has σ -o.c.-norm (that is, $\|f_n\|_E \downarrow_n 0$ for any sequence $(f_n)_{n=1}^\infty$ in E^+ satisfying $f_n \downarrow_n 0$). For a sequence $(f_n)_{n=1}^\infty$ in E^+ the condition $f_n \downarrow_n 0$ is equivalent with $f_n(x) \downarrow 0$ for μ -a.e. $x \in \Omega$. By way of example, for $1 \leq p < \infty$ the space $E = L^p(\mu)$ has o.c.-norm but, $L^\infty(\mu)$ does not have o.c.-norm.

Let $(E, \|\cdot\|_E)$ be a B.f.s. over (Ω, Σ, μ) with carrier Ω . The *Köthe dual* (or *associate space*) E^\times of E is defined by

$$E^\times = \{g \in L^0(\mu) : \|g\|_{E^\times} < \infty\},$$

where

$$\|g\|_{E^\times} = \sup \left\{ \int_\Omega |fg| d\mu : f \in E, \|f\|_E \leq 1 \right\}, \quad g \in L^0(\mu). \tag{2}$$

Then $(E^\times, \|\cdot\|_{E^\times})$ is a B.f.s. over (Ω, Σ, μ) with carrier equal to Ω ; see [22], Sections 68, 69 and also [23], p. 418. From the definition it is also clear that Hölder’s inequality holds, that is,

$$\left| \int_{\Omega} fg \, d\mu \right| \leq \int_{\Omega} |fg| \, d\mu \leq \|f\|_E \|g\|_{E^\times}, \quad f \in E, \quad g \in E^\times. \tag{3}$$

According to [22], Section 69, Theorem 1, we also have

$$\|g\|_{E^\times} = \sup \left\{ \left| \int_{\Omega} fg \, d\mu \right| : f \in E, \|f\|_E \leq 1 \right\}, \quad g \in E^\times. \tag{4}$$

For each $g \in E^\times$, define the linear functional $\varphi_g : E \rightarrow \mathbb{C}$ by setting

$$\varphi_g(f) = \int_{\Omega} fg \, d\mu, \quad f \in E.$$

It is easy to see that $\varphi_g \in E^*$ (the dual Banach space of E) and that $\|\varphi_g\|_{E^*} = \|g\|_{E^\times}$ for all $g \in E^\times$ (see (4)). Consequently, the map $g \mapsto \varphi_g$, for $g \in E^\times$, is an isometric isomorphism from E^\times into E^* . Via this map, E^\times may be identified with a closed subspace of E^* and we frequently denote the dual pairing of E and E^\times by $\langle \cdot, \cdot \rangle$, that is,

$$\langle f, g \rangle = \int_{\Omega} fg \, d\mu, \quad f \in E, \quad g \in E^\times. \tag{5}$$

Since the carrier of E^\times is equal to Ω (whenever the carrier of E is Ω), it follows that E^\times separates the points of E (but, in general, E^\times is not norming).

The functionals in E^* which are of the form φ_g for some $g \in E^\times$ may be identified with the band E_n^* in E^* of all order continuous functionals on E ; see [22], Section 69, Theorem 3, and also [23], Section 112. In particular, $E^* = E^\times$ if and only if E has o.c.-norm (cf. Theorem 2.4.2 in [13]).

A B.f.s. E over (Ω, Σ, μ) has the Fatou property if, for any net $0 \leq u_\alpha \uparrow_\alpha$ in E satisfying $\sup_\alpha \|u_\alpha\|_E < \infty$, there exists $u \in E$ such that $0 \leq u_\alpha \uparrow_\alpha u$ and $\|u_\alpha\|_E \uparrow_\alpha \|u\|_E$. In this case, sequences also suffice; cf. [23], Theorem 113.2. By a theorem of Halperin and Luxemburg (see [22], Section 71, Theorem 1), a B.f.s. E has the Fatou property if and only if $E = E^{\times\times}$ isometrically. We point out that in [2] the Fatou property is included in the definition of a B.f.s.

3. Translation invariant banach function spaces

Let G be an infinite compact abelian group and denote by μ normalized Haar measure on the Borel σ -algebra $\mathcal{B}(G)$. The space of all regular, complex Borel measures in G is denoted by $M(G)$. The group operation is denoted by $+$ and the identity element of G is denoted by 0 . Moreover, $\mu(-A) = \mu(A)$ for all $A \in \mathcal{B}(G)$. For the general theory of locally compact abelian groups we refer the reader, for instance, to [11,19]. We denote $L^0(\mu)$ by the more traditional notation $L^0(G)$ and the corresponding L^p -spaces by $L^p(G)$, $1 \leq p \leq \infty$. A B.f.s. over $(G, \mathcal{B}(G), \mu)$ is simply called a B.f.s. over G . For each $y \in G$, define the translation operator $\tau_y : L^0(G) \rightarrow L^0(G)$ by setting

$$(\tau_y f)(x) = f(x - y), \quad x \in G,$$

for all $f \in L^0(G)$.

Definition 3.1. A translation invariant B.f.s over G is a B.f.s. $E \subseteq L^0(G)$ such that $\tau_y f \in E$ with $\|\tau_y f\|_E = \|f\|_E$ for all $y \in G$, whenever $f \in E$.

Such B.f.s.’ include all spaces $L^p(G)$, $1 \leq p \leq \infty$, all Orlicz spaces, Lorentz spaces and Marcinkiewicz spaces over G . Actually, any rearrangement invariant B.f.s. over G is translation invariant. However, there exist plenty of translation invariant B.f.s.’ which are *not* rearrangement invariant. We present an example (see also [Example 3.11](#)).

Example 3.2. Let G_1 and G_2 be two infinite compact abelian groups (with normalized Haar measure μ_1 and μ_2 , respectively) and consider the product group $G = G_1 \times G_2$ (with normalized Haar measure $\mu = \mu_1 \times \mu_2$). Let $1 < p, q < \infty$. Define the function norm $\|\cdot\|_{p \times q}$ on $L^0(G)$ by

$$\|f\|_{p \times q} = \left(\int_{G_1} \left(\int_{G_2} |f(x, y)|^q d\mu_2(y) \right)^{p/q} d\mu_1(x) \right)^{1/p}, \quad f \in L^0(G),$$

and let

$$E_{p \times q} = \{f \in L^0(G) : \|f\|_{p \times q} < \infty\}.$$

Equipped with the norm $\|\cdot\|_{p \times q}$ the space $E_{p \times q}$ is a translation invariant B.f.s. over G with the Fatou property and o.c.-norm. It is readily verified that $E_{p \times q}$ is *not* rearrangement invariant whenever $p \neq q$.

The following observation may be intuitively clear.

Lemma 3.3. *Let $E \neq \{0\}$ be a translation invariant B.f.s. over G . Then the carrier of E is G .*

Proof. Let $B \in \mathcal{B}(G)$ with $\mu(B) > 0$ be given. We have to show that there exists $C \in \mathcal{B}(G)$ satisfying $\mu(C) > 0$, $C \subseteq B$ and $\chi_C \in E$. By assumption there is $0 < f \in E$. Hence, there exist $\varepsilon > 0$ and $A \in \mathcal{B}(G)$ satisfying $\mu(A) > 0$ and $0 < \varepsilon \chi_A \leq f$. In particular, $\chi_A \in E$. It follows from Theorem F in Section 59 of [10] that there exists $y_0 \in G$ such that $(\tau_{y_0} \chi_A) \chi_B > 0$. Setting $C = (A + y_0) \cap B$, it follows that $\mu(C) > 0$ and $\chi_C \leq \tau_{y_0} \chi_A \in E$, which implies that $\chi_C \in E$. Hence, the set C satisfies the requirements and the proof is complete. ■

Using the definition (see (2)), it is routine to verify that the Köthe dual E^\times is translation invariant whenever $E \neq \{0\}$ is a translation invariant B.f.s. over G . If $E \neq \{0\}$ is any rearrangement invariant B.f.s. over G , then $L^\infty(G) \subseteq E \subseteq L^1(G)$ (as μ is atomless with $\mu(G) < \infty$). The following result shows, for any translation invariant B.f.s. E over G , that the inclusion $E \subseteq L^1(G)$ always holds. Even if $E \neq \{0\}$, the other inclusion $L^\infty(G) \subseteq E$ may fail, in general (see [Example 3.6](#)).

Proposition 3.4. *Let E be any translation invariant B.f.s. over G . Then $E \subseteq L^1(G)$ with a continuous embedding.*

Proof. We can assume that $E \neq \{0\}$. Since then also $E^\times \neq \{0\}$, there exists $0 \leq g \in E^\times$ with $\|g\|_{E^\times} = 1$. Given any $0 \leq f \in E$, the non-negative function $(x, y) \mapsto f(x + y)g(y)$, for $(x, y) \in G \times G$, is known to be Borel measurable on $G \times G$ ([19], p. 5). Since $\int_G f d\mu = \int_G \tau_{-y} f d\mu$ for all $y \in G$ (the value $+\infty$ is possible at this stage), we deduce

from the Fubini–Tonelli theorem (for positive functions) that

$$\begin{aligned} & \int_G \left(\int_G f(x+y) g(y) d\mu(y) \right) d\mu(x) \\ &= \int_G \left(\int_G f(x+y) d\mu(x) \right) g(y) d\mu(y) \\ &= \left(\int_G f d\mu \right) \left(\int_G g d\mu \right). \end{aligned}$$

But, for each $x \in G$, it follows from (3) that

$$\int_G f(x+y) g(y) d\mu(y) = \int_G (\tau_{-y}f) g d\mu \leq \|f\|_E \|g\|_{E^\times} = \|f\|_E < \infty$$

and so, by the previous identity, we have that

$$\left(\int_G f d\mu \right) \left(\int_G g d\mu \right) \leq \mu(G) \|f\|_E = \|f\|_E < \infty. \tag{6}$$

Since $g > 0$, its integral $\alpha = \int_G g d\mu > 0$ and so $\|f\|_1 = \int_G f d\mu < \infty$, that is, $f \in L^1(G)$. It follows that $E \subseteq L^1(G)$. As noted earlier, the natural embedding $E \subseteq L^1(G)$ is necessarily continuous. ■

Corollary 3.5. *Let $E \neq \{0\}$ be a translation invariant B.f.s. over G .*

- (i) *With continuous inclusions we have that $L^\infty(G) \subseteq E^\times \subseteq L^1(G)$.*
- (ii) *If, in addition, E has the Fatou property, then $L^\infty(G) \subseteq E \subseteq L^1(G)$, with continuous inclusions.*

Proof. (i). Since $E^\times \neq \{0\}$ is also a translation invariant B.f.s., we can apply Proposition 3.4 to E^\times to conclude that $E^\times \subseteq L^1(G)$ continuously. Again by Proposition 3.4, now applied to E , we have $E \subseteq L^1(G)$ continuously and hence, $L^\infty(G) = L^1(G)^\times \subseteq E^\times$, continuously.

(ii). Part (i), applied to E^\times , yields $L^\infty(G) \subseteq E^{\times\times} \subseteq L^1(G)$ continuously. If E has the Fatou property, then $E = E^{\times\times}$, which completes the proof. ■

There exist non-trivial translation invariant B.f.s.’ over G for which $L^\infty(G) \not\subseteq E$.

Example 3.6. This example is inspired by the results in [20]. Let G be an infinite compact abelian group, equipped with normalized Haar measure μ . Let \mathcal{O} denote the collection of all open and dense subsets of G . Note that $U_1 \cap U_2 \in \mathcal{O}$ whenever $U_1, U_2 \in \mathcal{O}$ and that $x + U \in \mathcal{O}$ whenever $U \in \mathcal{O}$ and $x \in G$. Define

$$J_{\mathcal{O}} = \{f \in L^\infty(G) : \exists U \in \mathcal{O} \text{ such that } f\chi_U = 0\}.$$

It is readily verified that $J_{\mathcal{O}}$ is an ideal in $L^\infty(G)$, which is also translation invariant (that is, $\tau_x f \in J_{\mathcal{O}}$ whenever $f \in J_{\mathcal{O}}$ and $x \in G$). Furthermore, $J_{\mathcal{O}} \neq \{0\}$. Indeed, there exists $U \in \mathcal{O}$ with $\mu(U) < 1$ (actually, for each $\varepsilon > 0$ there exists $V \in \mathcal{O}$ such that $\mu(V) < \varepsilon$; see Theorem 2.4 in [20]), in which case $0 \neq \chi_{G \setminus U} \in J_{\mathcal{O}}$. Let $E_{\mathcal{O}}$ denote the norm closure of $J_{\mathcal{O}}$ in $L^\infty(G)$. Since the norm closure of any ideal is again an ideal (cf. Theorem 100.11 in [23]), it follows that $E_{\mathcal{O}}$ is an ideal in $L^\infty(G)$ and it is easily verified that $E_{\mathcal{O}}$ is translation invariant. Hence, $(E_{\mathcal{O}}, \|\cdot\|_\infty)$ is a non-zero translation invariant B.f.s. over G .

The claim is that $\chi_G \notin E_{\mathcal{O}}$ (and hence, that $L^\infty(G) \not\subseteq E_{\mathcal{O}}$). Actually, $E_{\mathcal{O}} \cap C(G) = \{0\}$. Indeed, suppose that $0 \neq f \in E_{\mathcal{O}} \cap C(G)$. Then there exist an $\varepsilon > 0$ and a non-empty open

set $W \subseteq G$ such that $|f(x)| > \varepsilon$, for $x \in W$. Choose $g \in J_{\mathcal{O}}$ such that $\|f - g\|_{\infty} < \varepsilon/2$, in which case $|g(x)| > \varepsilon/2$ for all $x \in W$. But, then g cannot vanish μ -a.e. on any set from \mathcal{O} , contradicting that $g \in J_{\mathcal{O}}$.

Our next aim is to show that $L^{\infty}(G) \subseteq E$ whenever $E \neq \{0\}$ and E has o.c.-norm (Proposition 3.8). First we require some preliminary results.

Lemma 3.7. *Let $E \neq \{0\}$ be a translation invariant B.f.s. over G with o.c.-norm. For each set $A \in \Sigma$ with $\chi_A \in E$ the following assertions hold.*

- (i) *For each $\varepsilon > 0$ there exists $\delta > 0$ such that $\|\chi_B\|_E \leq \varepsilon$ whenever $B \in \mathcal{B}(G)$ satisfies $B \subseteq A$ and $\mu(B) \leq \delta$.*
- (ii) *If a sequence $(A_n)_{n=1}^{\infty}$ in $\mathcal{B}(G)$ satisfies $A_n \downarrow \emptyset$, then*

$$\limsup_{n \rightarrow \infty} \sup_{y \in G} \|\chi_{A_n}(\tau_y \chi_A)\|_E = 0.$$

- (iii) *For each $B \in \mathcal{B}(G)$ we have that $\lim_{y \rightarrow 0} \|(\chi_B - \tau_y \chi_B) \chi_A\|_E = 0$.*
- (iv) $\lim_{y \rightarrow 0} \|\chi_A - \tau_y \chi_A\|_E = 0$.

Proof. (i). If the statement does not hold, then there exist $\varepsilon > 0$ and a sequence $(B_n)_{n=1}^{\infty}$ in $\mathcal{B}(G)$ with $B_n \subseteq A$, $\mu(B_n) \leq 2^{-n}$ and $\|\chi_{B_n}\|_E \geq \varepsilon$ for all $n \in \mathbb{N}$. The sets $C_n = \bigcup_{k=n}^{\infty} B_k$ satisfy $\mu(C_n) \leq 2^{-n+1}$ for all $n \in \mathbb{N}$ and $C_n \downarrow \emptyset$. Hence, $\chi_A \geq \chi_{C_n} \downarrow 0$ in E and so, by order continuity of the norm, we have $\|\chi_{C_n}\|_E \downarrow 0$. But, $\chi_{C_n} \geq \chi_{B_n}$ implies that $\|\chi_{C_n}\|_E \geq \|\chi_{B_n}\|_E \geq \varepsilon$ for all n , which is a contradiction.

(ii). Given $\varepsilon > 0$, by part (i) there exists $\delta > 0$ such that $\|\chi_B\|_E \leq \varepsilon$ whenever $B \in \mathcal{B}(G)$ satisfies $B \subseteq A$ and $\mu(B) \leq \delta$. Since $A_n \downarrow \emptyset$, there exists $N \in \mathbb{N}$ such that $\mu(A_n) \leq \delta$ for all $n \geq N$. Observe that

$$\|\chi_{A_n}(\tau_y \chi_A)\|_E = \|\tau_{-y}(\chi_{A_n} \tau_y \chi_A)\|_E = \|(\tau_{-y} \chi_{A_n}) \chi_A\|_E, \quad n \in \mathbb{N}, \quad y \in G.$$

Since $(\tau_{-y} \chi_{A_n}) \chi_A = \chi_{(A_n - y) \cap A}$ and

$$\mu((A_n - y) \cap A) \leq \mu(A_n - y) = \mu(A_n) \leq \delta, \quad n \geq N,$$

it follows that $\|(\tau_{-y} \chi_{A_n}) \chi_A\|_E \leq \varepsilon$ for all $y \in G$ and $n \geq N$. Hence,

$$\sup_{y \in G} \|(\tau_{-y} \chi_{A_n}) \chi_A\|_E \leq \varepsilon, \quad n \geq N,$$

which completes the proof of part (ii).

(iii). Fix $B \in \mathcal{B}(G)$. Note that $|\chi_B - \tau_y \chi_B| = \chi_{(B+y) \Delta B}$, for $y \in G$, where Δ denotes the symmetric difference. It follows from the continuity of translations in $L^1(G)$ (see [19], p.3) that $\lim_{y \rightarrow 0} \|\chi_B - \tau_y \chi_B\|_1 = 0$ and hence, that $\lim_{y \rightarrow 0} \mu((B+y) \Delta B) = 0$. Given $\varepsilon > 0$, it follows from part (i) that there exists $\delta > 0$ such that $\|\chi_C\|_E \leq \varepsilon$ whenever $C \in \mathcal{B}(G)$ satisfies $C \subseteq A$ and $\mu(C) \leq \delta$. If U is a neighbourhood of $0 \in G$ such that $\mu((B+y) \Delta B) \leq \delta$ for all $y \in U$, then $\mu(A \cap ((B+y) \Delta B)) \leq \delta$ and so,

$$\|(\chi_B - \tau_y \chi_B) \chi_A\|_E = \|\chi_{A \cap ((B+y) \Delta B)}\|_E \leq \varepsilon, \quad y \in U.$$

This suffices for the proof of part (iii).

(iv). Since the carrier of E is G (cf. Lemma 3.3), there exists a sequence $(H_n)_{n=1}^{\infty}$ in $\mathcal{B}(G)$ with $\chi_{H_n} \in E$ for all n and $H_n \uparrow G$, i.e., $G \setminus H_n \downarrow \emptyset$. Given $\varepsilon > 0$, it follows from part (ii) that there is $N \in \mathbb{N}$ such that $\sup_{y \in G} \|\chi_{G \setminus H_n}(\tau_y \chi_A)\|_E \leq \varepsilon/3$. By part (iii), there is a

neighbourhood U of $0 \in G$ such that $\|\chi_{H_N}(\chi_A - \tau_y \chi_A)\|_E \leq \varepsilon/3$ for all $y \in U$. Consequently, for each $y \in U$ we have

$$\begin{aligned} \|\chi_A - \tau_y \chi_A\|_E &\leq \|\chi_{H_N}(\chi_A - \tau_y \chi_A)\|_E + \|\chi_{G \setminus H_N}(\chi_A - \tau_y \chi_A)\|_E \\ &\leq \varepsilon/3 + \|\chi_{G \setminus H_N}(\tau_0 \chi_A)\|_E + \|\chi_{G \setminus H_N}(\tau_y \chi_A)\|_E \leq \varepsilon. \end{aligned}$$

This completes the proof of part (iv). ■

We can now establish the result alluded to above.

Proposition 3.8. *Let $E \neq \{0\}$ be a translation invariant B.f.s. over G with o.c.-norm. The following statements hold.*

- (i) For each $f \in E$ it is the case that $\lim_{y \rightarrow 0} \|f - \tau_y f\|_E = 0$.
- (ii) $L^\infty(G) \subseteq E \subseteq L^1(G)$ with continuous inclusions.
- (iii) $C(G)$ is a dense subspace of E .

Proof. (i). Denote by $\text{sim}(\mathcal{B}(G))$ the space of all simple functions based on $\mathcal{B}(G)$. Claim: $\text{sim}(\mathcal{B}(G)) \cap E$ is dense in E . Indeed, given $0 \leq f \in E$, there exists a sequence $(s_n)_{n=1}^\infty$ in $\text{sim}(\mathcal{B}(G))^+$ such that $0 \leq s_n \uparrow_n f$ μ -a.e. on G . Since E is an ideal in $L^0(G)$, it is clear that $s_n \in \text{sim}(\mathcal{B}(G)) \cap E$, for $n \in \mathbb{N}$. The order continuity of the norm in E implies that $\|f - s_n\|_E \rightarrow 0$ as $n \rightarrow \infty$. This suffices for the proof of the claim.

Let $s \in \text{sim}(\mathcal{B}(G)) \cap E$. Then $s = \sum_{j=1}^k \alpha_j \chi_{A_j}$, where $A_j \in \mathcal{B}(G)$ with $\chi_{A_j} \in E$ and $\alpha_j \in \mathbb{C}$, for $j = 1, \dots, k$. Therefore, Lemma 3.7 (iv) implies that $\|s - \tau_y s\|_E \rightarrow 0$ as $y \rightarrow 0$ in G . Since $\text{sim}(\mathcal{B}(G)) \cap E$ is dense in E , the result of (i) is now clear.

(ii). That $E \subseteq L^1(G)$ has been shown in Proposition 3.4. For the inclusion of $L^\infty(G)$ into E , we begin with a general observation, which is of independent interest (only special cases will be used in the proofs of (ii) and (iii)).

Let $f \in E$ and $\lambda \in M(G)$. Since $f \in L^1(G)$ (cf. Proposition 3.4), the convolution $f * \lambda \in L^1(G)$ exists (see e.g. [19], Section 1.3.2). The claim is that $f * \lambda \in E$. Indeed, it follows from part (i) that the function $F : y \mapsto \tau_y f$, for $y \in G$, is continuous from G into E . So, the range of F is a compact metric space and hence, is separable. Via the Pettis measurability theorem ([6], p. 42) it follows that F is strongly $|\lambda|$ -measurable and bounded. Consequently, the Bochner integral $\int_G^{(B)} F d\lambda = \int_G^{(B)} \tau_y f d\lambda(y) \in E \subseteq L^1(G)$ exists. Using [6], Theorem II.2.6 (for continuous functionals on E) and Fubini’s theorem, we find for all $g \in L^\infty(G) \subseteq E^\times \subseteq E^*$ that

$$\begin{aligned} \left\langle \int_G^{(B)} \tau_y f d\lambda(y), g \right\rangle &= \int_G \langle \tau_y f, g \rangle d\lambda(y) \\ &= \int_G \left(\int_G (\tau_y f)(x) g(x) d\mu(x) \right) d\lambda(y) \\ &= \int_G \left(\int_G f(x-y) d\lambda(y) \right) g(x) d\mu(x) = \langle f * \lambda, g \rangle. \end{aligned}$$

We can conclude that

$$f * \lambda = \int_G^{(B)} \tau_y f d\lambda(y) \in E, \quad f \in E, \quad \lambda \in M(G). \tag{7}$$

This proves the claim.

Fix any $0 < f_0 \in E$. By what has just been proved, we have that $f_0 * \mu \in E$. But, $f_0 * \mu = (\int_G f_0 d\mu) \chi_G$ and so we can conclude that $\chi_G \in E$ and hence, that $L^\infty(G) \subseteq E$. Furthermore, if $f \in L^\infty(G)$, then $|f| \leq \|f\|_\infty \chi_G$ implies that $\|f\|_E \leq \|\chi_G\|_E \|f\|_\infty$. Consequently, the embedding $L^\infty(G) \subseteq E$ is continuous.

(iii). Let $f \in E$ be fixed and $\varepsilon > 0$ be given. It follows from (i) that there exists an open neighbourhood U of 0 in G such that $\|f - \tau_y f\|_E \leq \varepsilon$ for all $y \in U$. Defining $h = \mu(U)^{-1} \chi_U$, we have $f * h \in C(G)$ (as $f \in L^1(G)$ and $h \in L^\infty(G)$; cf. [19], Section 1.1.6). Furthermore, (7) implies that

$$f - (f * h) = \mu(U)^{-1} \int_U^{(B)} (f - \tau_y f) d\mu(y),$$

which implies that

$$\|f - (f * h)\|_E \leq \mu(U)^{-1} \int_U \|f - \tau_y f\|_E d\mu(y) \leq \varepsilon;$$

cf. [6], Theorem II.2.4 (ii). The proof is thereby complete. ■

Remark 3.9.

- (a) Let $E \neq \{0\}$ be a translation invariant B.f.s. over G with o.c.-norm. Then $f * \lambda \in E$ for all $f \in E$ and $\lambda \in M(G)$; see the proof of part (ii) in the above proposition. Moreover, it follows from (7) that $\|f * \lambda\|_E \leq \|\lambda\|_{M(G)} \|f\|_E$. Consequently, for each $\lambda \in M(G)$, the operator $T_\lambda^E : E \rightarrow E$ of convolution with λ exists and satisfies $\|T_\lambda^E\| \leq \|\lambda\|_{M(G)}$. It can be shown that a similar result holds for translation invariant B.f.s.’ over G which have the Fatou property. However, for general translation invariant B.f.s.’ over G this result fails. In fact, for the translation invariant B.f.s. $E_{\mathcal{O}}$ of Example 3.6 it can be shown that the only measures $\lambda \in M(G)$ for which the convolution operator $T_\lambda^{E_{\mathcal{O}}} : E_{\mathcal{O}} \rightarrow E_{\mathcal{O}}$ exists, are the discrete measures. See [15].
- (b) A translation invariant B.f.s. E over G with the property that $\|f - \tau_y f\|_E \rightarrow 0$ as $y \rightarrow 0$ in G , for every $f \in E$, necessarily has o.c.-norm. The proof of this fact is somewhat involved and is not needed in this note. See [15]. This fact also implies that any translation invariant B.f.s. over G in which $C(G)$ is dense, necessarily has o.c.-norm.

Next we discuss the phenomenon that a general translation invariant B.f.s. need not be reflection invariant (see the definition below). For any $f \in L^0(G)$, its reflection $\tilde{f} \in L^0(G)$ is defined by

$$\tilde{f}(x) = f(-x), \quad x \in G.$$

Definition 3.10. A B.f.s. E over G is called *reflection invariant* if it has the property that $\tilde{f} \in E$ with $\|\tilde{f}\|_E = \|f\|_E$ whenever $f \in E$.

Evidently, every rearrangement invariant B.f.s. over G is reflection invariant. Note that the B.f.s. $E_{p \times q}$ of Example 3.2, with $p \neq q$, is translation and reflection invariant but, it is not rearrangement invariant. Furthermore, it is routine to verify, for any reflection invariant B.f.s. E over G with carrier equal to G , that the Köthe dual E^\times is also reflection invariant. However, a translation invariant B.f.s. over G need not be reflection invariant.

Example 3.11. Let $0 < g \in L^1(G)$ be fixed. Define

$$\|f\|_{E_g} = \sup_{y \in G} \int_G |f|(\tau_y g) d\mu, \quad f \in L^0(G), \tag{8}$$

and let

$$E_g = \left\{ f \in L^0(G) : \|f\|_{E_g} < \infty \right\}, \tag{9}$$

equipped with the norm $\|\cdot\|_{E_g}$. It is readily verified that E_g is a translation invariant B.f.s. over G with the Fatou property. Actually, E_g is the largest translation invariant B.f.s. F over G with the property that $g \in F^\times$.

Consider the group $G = \mathbb{T}$ with normalized Haar measure (that is, $d\mu = \frac{1}{2\pi}dx$ on $[-\pi, \pi]$, where dx denotes Lebesgue measure). Functions on \mathbb{T} will be identified with 2π -periodic functions on \mathbb{R} . The 2π -periodic function $g : \mathbb{R} \rightarrow \mathbb{R}$ is defined by setting

$$g(x) = \begin{cases} 0 & \text{if } -\pi < x \leq 0 \\ \frac{1}{\sqrt{x}} & \text{if } 0 < x \leq \pi \end{cases}.$$

Define the function norm $\|\cdot\|_{E_g}$ on $L^0(G)$ by (8) and let E_g be the translation invariant B.f.s. defined by (9). It is clear that the function $g \notin E_g$. However, a direct computation shows that $\tilde{g} \in E_g$. Therefore, E_g is not reflection invariant. It can be shown that the space E_g does not have o.c.-norm. There also exist translation invariant B.f.s.' over \mathbb{T} which are not reflection invariant but, have both the Fatou property and o.c.-norm.

4. Fourier multiplier functions

As before, G is an infinite compact abelian group equipped with normalized Haar measure μ . Let Γ be the dual group of G ; see, for example, Section 1.2 of [19]. For $x \in G$ and $\gamma \in \Gamma$ we write $\gamma(x) = (x, \gamma)$. Since we frequently consider Γ as a subset of $C(G)$, we shall write the group operation in Γ as *multiplication*, that is, if $\gamma_1, \gamma_2 \in \Gamma$, then $\gamma_1\gamma_2 \in \Gamma$ is given by

$$(x, \gamma_1\gamma_2) = (x, \gamma_1)(x, \gamma_2), \quad x \in G.$$

The identity element of Γ is, of course, χ_G . In particular,

$$(x, \gamma^{-1}) = \overline{(x, \gamma)} = (-x, \gamma), \quad x \in G, \quad \gamma \in \Gamma, \tag{10}$$

where the bar denotes complex conjugation. Denote by $\tau(G)$ the linear subspace of $C(G)$ consisting of all *trigonometric polynomials* on G , that is, $\tau(G)$ is the linear span of $\Gamma \subseteq C(G)$. The following fact, which is a consequence of Proposition 3.8 (iii), will be used.

Proposition 4.1. *Let $E \neq \{0\}$ be a translation invariant B.f.s. over G with o.c.-norm. Then $\tau(G)$ is a dense subspace of E .*

Proof. The dual group Γ separates the points of G (see [19], Section 1.5.2) and so it follows from the Stone–Weierstrass theorem that $\tau(G)$ is dense in $C(G)$ with respect to $\|\cdot\|_\infty$. By Proposition 3.8 (iii), $C(G)$ is dense in E . Since the embedding of $C(G)$ into E is continuous (as $L^\infty(G) \subseteq E$ continuously), we can conclude that $\tau(G)$ is dense in E . ■

For each $f \in L^1(G)$ its Fourier transform, denoted by $\hat{f} : \Gamma \rightarrow \mathbb{C}$, is given by

$$\hat{f}(\gamma) = \int_G f(x)(-x, \gamma) d\mu(x), \quad \gamma \in \Gamma,$$

and satisfies $\hat{f} \in c_0(\Gamma)$ with $\|\hat{f}\|_\infty \leq \|f\|_1$. Furthermore, if $f \in L^1(G)$ and $\hat{f} = 0$, then $f = 0$. For these facts, see for instance Section 1.2 of [19].

Let $E \neq \{0\}$ be translation invariant B.f.s.’ over G . Recall that E is contained in $L^1(G)$ and so \hat{f} exists for $f \in E$. A function $\varphi : \Gamma \rightarrow \mathbb{C}$ is called an E -multiplier function for G if, for every $f \in E$, there exists a function $g \in E$ such that $\hat{g} = \varphi \hat{f}$ (pointwise on Γ). If such a function g exists, then it is necessarily unique by the injectivity of the Fourier transform. Denote g by $M_\varphi^E f$. The so defined map $M_\varphi^E : E \rightarrow E$ is linear and satisfies

$$(M_\varphi^E f)^\wedge = \varphi \hat{f}, \quad f \in E. \tag{11}$$

The collection of all E -multiplier functions for G is denoted by $\mathcal{M}_E(G)$, which clearly is a (complex) vector space of functions on Γ . For $E = L^p(G)$, where $1 \leq p \leq \infty$, we use the simpler notation $\mathcal{M}_p(G)$ in place of $\mathcal{M}_{L^p(G)}(G)$. A routine application of the closed graph theorem shows that M_φ^E is continuous, i.e., $M_\varphi^E \in \mathcal{L}(E)$, for every $\varphi \in \mathcal{M}_E(G)$. Here, $\mathcal{L}(E)$ denotes the Banach space of all continuous linear operators of E into itself, equipped with the operator norm. It should be observed that if E is a translation invariant B.f.s. over G with $L^\infty(G) \subseteq E$ (in which case $\gamma \in E$ for all $\gamma \in \Gamma$) then, for all $\varphi \in \mathcal{M}_E(G)$, we have

$$M_\varphi^E \gamma = \varphi(\gamma) \gamma, \quad \gamma \in \Gamma. \tag{12}$$

Indeed, given $\varphi \in \mathcal{M}_E(G)$ and $\gamma \in \Gamma$, we have that $\hat{\gamma} = \chi_{\{\gamma\}}$, from which it follows that $(M_\varphi^E \gamma)^\wedge = \varphi \hat{\gamma} = (\varphi(\gamma) \gamma)^\wedge$. This implies (12). It follows from (11) that $\mathcal{M}_E(G)$ is an algebra for pointwise multiplication of functions on Γ .

It is shown in the following lemma that always $\mathcal{M}_E(G) \subseteq \ell^\infty(\Gamma)$. Recall that if $E \neq \{0\}$ is a translation invariant B.f.s. over G , then $L^\infty(G) \subseteq E^\times$; Corollary 3.5(i). In particular, $\gamma \in E^\times$ for each $\gamma \in \Gamma$. Moreover, $\|\gamma\|_{E^\times} = \|\chi_G\|_{E^\times}$ for all $\gamma \in \Gamma$, because $|\gamma| = \chi_G$. Recall also that $\|f\|_{E^\times} = \|f\|_{E^*}$ for each $f \in E^\times$; see the discussion following formula (4).

Lemma 4.2. *Let $E \neq \{0\}$ be a translation invariant B.f.s.’ over G . Then $\mathcal{M}_E(G) \subseteq \ell^\infty(\Gamma)$ and*

$$\|\varphi\|_\infty \leq \|M_\varphi^E\|, \quad \varphi \in \mathcal{M}_E(G). \tag{13}$$

Proof. Fix $\varphi \in \mathcal{M}_E(G)$ and set $T = M_\varphi^E$. As observed prior to the lemma, T is continuous and so we can consider the Banach space adjoint operator $T^* : E^* \rightarrow E^*$. Let $\gamma \in \Gamma$ and consider $\gamma \in E^\times \subseteq E^*$. The claim is that $T^* \gamma = \varphi(\gamma^{-1}) \gamma \in E^\times \subseteq E^*$. Indeed, given $f \in E$ we have that

$$\begin{aligned} \langle f, T^* \gamma \rangle &= \langle Tf, \gamma \rangle = \int_G (Tf)(x) (x, \gamma) d\mu(x) \\ &= \int_G (Tf)(x) (-x, \gamma^{-1}) d\mu(x) = (Tf)^\wedge(\gamma^{-1}) \\ &= \varphi(\gamma^{-1}) \hat{f}(\gamma^{-1}) = \varphi(\gamma^{-1}) \int_G f(x) (-x, \gamma^{-1}) d\mu(x) \\ &= \varphi(\gamma^{-1}) \int_G f(x) (x, \gamma) d\mu(x) = \langle f, \varphi(\gamma^{-1}) \gamma \rangle. \end{aligned}$$

Since E separates the elements of E^* , the claim follows. Defining the *reflected function* $\tilde{\varphi} : \Gamma \rightarrow \mathbb{C}$ by $\tilde{\varphi}(\gamma) = \varphi(\gamma^{-1})$, for $\gamma \in \Gamma$, we find that

$$\|T\| = \|T^*\| \geq \sup_{\gamma \in \Gamma} \frac{\|T^* \gamma\|_{E^\times}}{\|\gamma\|_{E^\times}} = \sup_{\gamma \in \Gamma} \frac{|\tilde{\varphi}(\gamma)| \|\gamma\|_{E^\times}}{\|\gamma\|_{E^\times}} = \|\tilde{\varphi}\|_\infty.$$

Since $\|\tilde{\varphi}\|_\infty = \|\varphi\|_\infty$, this suffices for the proof. ■

For the case $E = L^p(G)$ with $1 \leq p < \infty$, the inequality (13) is well known; see Corollary 4.1.2 of [11], for example.

Define the family of operators

$$\mathcal{M}_E^{\text{op}}(G) = \{M_\varphi^E : \varphi \in \mathcal{M}_E(G)\} \subseteq \mathcal{L}(E). \tag{14}$$

For each $x \in G$, a direct calculation shows that the Dirac point measure δ_x satisfies $\tau_x f = f * \delta_x$, for $f \in E$. Since $\tau_x \in \mathcal{L}(E)$ and $\hat{\delta}_x(\gamma) = (-x, \gamma)$, for $\gamma \in \Gamma$, it follows that $(x, \cdot) \in \mathcal{M}_E(G)$ for every $x \in G$ and $\{\tau_x : x \in G\} \subseteq \mathcal{M}_E^{\text{op}}(G)$. In particular, the identity operator $I = \tau_0 \in \mathcal{M}_E^{\text{op}}(G)$ and the constant function $\hat{\delta}_0 = \mathbf{1} \in \mathcal{M}_E(G)$.

The following result is a consequence of Lemma 4.2.

Corollary 4.3. *Let $E \neq \{0\}$ be a translation invariant B.f.s. over G . The set $\mathcal{M}_E(G)$ is a unital subalgebra of $\ell^\infty(\Gamma)$ and the map $\Phi : \mathcal{M}_E(G) \rightarrow \mathcal{L}(E)$, defined by*

$$\Phi(\varphi) = M_\varphi^E, \quad \varphi \in \mathcal{M}_E(G),$$

is a unital algebra isomorphism onto the subalgebra $\mathcal{M}_E^{\text{op}}(G) \subseteq \mathcal{L}(E)$. Moreover, $\mathcal{M}_E^{\text{op}}(G)$ is a unital commutative Banach subalgebra of $\mathcal{L}(E)$.

Proof. It is routine to verify that $\mathcal{M}_E(G)$ is a unital subalgebra of $\ell^\infty(\Gamma)$ and that the map $\Phi : \mathcal{M}_E(G) \rightarrow \mathcal{L}(E)$ is a unital algebra homomorphism. Consequently, $\mathcal{M}_E^{\text{op}}(G) = \Phi(\mathcal{M}_E(G))$ is a unital commutative subalgebra of $\mathcal{L}(E)$. The injectivity of Φ follows from Lemma 4.2.

To show that $\mathcal{M}_E^{\text{op}}(G)$ is norm closed in $\mathcal{L}(E)$, let $(M_{\varphi_n}^E)_{n=1}^\infty$ be a sequence in $\mathcal{M}_E^{\text{op}}(G)$ satisfying $\|S - M_{\varphi_n}^E\| \rightarrow 0$ as $n \rightarrow \infty$ for some $S \in \mathcal{L}(E)$. Lemma 4.2 implies that $(\varphi_n)_{n=1}^\infty$ is a Cauchy sequence in $\ell^\infty(\Gamma)$. So, there exists $\varphi \in \ell^\infty(\Gamma)$ such that $\|\varphi - \varphi_n\|_\infty \rightarrow 0$ for $n \rightarrow \infty$. It follows, for a fixed $f \in E \subseteq L^1(G)$, that $\hat{f} \in c_0(\Gamma)$ and so

$$(M_{\varphi_n}^E f)^\wedge = \varphi_n \hat{f} \rightarrow \varphi \hat{f}, \quad n \rightarrow \infty, \tag{15}$$

in $\ell^\infty(\Gamma)$. On the other hand, $M_{\varphi_n}^E f \rightarrow S f$ in E for $n \rightarrow \infty$. Since E is continuously included in $L^1(G)$, it follows that $M_{\varphi_n}^E f \rightarrow S f$ in $L^1(G)$. By the continuity of the Fourier transform from $L^1(G)$ into $c_0(\Gamma)$ we can conclude that $(M_{\varphi_n}^E f)^\wedge \rightarrow (S f)^\wedge$ in $\ell^\infty(\Gamma)$. Then (15) implies that $(S f)^\wedge = \varphi \hat{f}$. But, $f \in E$ is arbitrary and so $\varphi \in \mathcal{M}_E(G)$ with $S = M_\varphi^E$. The proof is complete. ■

Remark 4.4.

- (a) For $E = L^2(G)$ it is a classical result that $\mathcal{M}_2(G) = \ell^\infty(\Gamma)$, which is an immediate consequence of the fact that the Fourier transform is a surjective isometry from $L^2(G)$ onto $\ell^2(\Gamma)$. Conversely, if E is a translation invariant B.f.s. over G with $L^\infty(G) \subseteq E$ and $\mathcal{M}_E(G) = \ell^\infty(\Gamma)$, then it follows from Proposition 1.6 of [5] that $E = L^2(G)$ with equivalent norms.
- (b) Let $E \neq \{0\}$ be a translation invariant B.f.s. over G with *o.c.-norm*. As observed in Remark 3.9, for each $\lambda \in M(G)$ the operator $T_\lambda^E : E \rightarrow E$ of convolution with λ exists. Recalling that $(f * \lambda)^\wedge = \hat{\lambda} \hat{f}$ for $f \in L^1(G)$ and $\lambda \in M(G)$, it follows that $\hat{\lambda} \in \mathcal{M}_E(G)$ with $M_{\hat{\lambda}}^E = T_\lambda^E$ for all $\lambda \in M(G)$. Hence,

$$\{\hat{\lambda} : \lambda \in M(G)\} \subseteq \mathcal{M}_E(G).$$

A similar result holds if E has the Fatou property.

For any function $f \in L^0(G)$, the function $f^\natural \in L^0(G)$ is defined by

$$f^\natural(x) = \overline{f(-x)}, \quad \mu\text{-a.e } x \in G.$$

Let E be a reflection and translation invariant B.f.s. over G . Then

$$f^\natural \in E \quad \text{and} \quad \|f^\natural\|_E = \|f\|_E, \quad f \in E,$$

and the map $f \mapsto f^\natural$, for $f \in E$, is a conjugate linear isometric involution in E . For each operator $T \in \mathcal{L}(E)$ the operator $T^\natural \in \mathcal{L}(E)$ is defined by

$$T^\natural f = (Tf)^\natural, \quad f \in E,$$

and satisfies $\|T^\natural\| = \|T\|$. Then $T \mapsto T^\natural$, for $T \in \mathcal{L}(E)$, is a conjugate linear isometric involution in $\mathcal{L}(E)$.

Proposition 4.5. *Let $E \neq \{0\}$ be a translation and reflection invariant B.f.s. over G .*

- (i) *If $\varphi \in \mathcal{M}_E(G)$, then also $\bar{\varphi} \in \mathcal{M}_E(G)$ and $M_{\bar{\varphi}}^E = (M_\varphi^E)^\natural$.*
- (ii) *The map $M_\varphi^E \mapsto M_{\bar{\varphi}}^E$, for $\varphi \in \mathcal{M}_E(G)$, is a conjugate linear isometric involution in $\mathcal{M}_E^{\text{op}}(G)$.*

Proof. (i). Let $\varphi \in \mathcal{M}_E(G)$. A direct (but careful) computation shows that

$$\left((M_\varphi^E)^\natural f \right)^\wedge = \bar{\varphi} \hat{f}, \quad f \in E.$$

Consequently, $\bar{\varphi} \in \mathcal{M}_E(G)$ and $M_{\bar{\varphi}}^E = (M_\varphi^E)^\natural$.

(ii). This follows immediately from part (i) in combination with the observations made prior to the present proposition. ■

For $E = L^2(G)$ and $\varphi \in \mathcal{M}_2(G) = \ell^\infty(\Gamma)$, it follows (via Plancherel’s formula) that $M_{\bar{\varphi}}^{(2)} = (M_\varphi^{(2)})^*$ is the Hilbert space adjoint of $M_\varphi^{(2)}$.

5. A Fuglede type theorem

Again let G be an infinite compact abelian group with normalized Haar measure μ . As observed in Proposition 4.5(i), if E is a translation and reflection invariant B.f.s. over G , then $\bar{\varphi} \in \mathcal{M}_E(G)$ whenever $\varphi \in \mathcal{M}_E(G)$. For $E = L^2(G)$, we noted that $M_{\bar{\varphi}}^{(2)} = (M_\varphi^{(2)})^*$ is the Hilbert space adjoint of $M_\varphi^{(2)}$. In particular, $M_\varphi^{(2)}$ is a normal operator on $L^2(G)$ (as $\mathcal{M}_2^{\text{op}}(G)$ is commutative; cf. Corollary 4.3). Consequently, it follows from Fuglede’s theorem (see, for instance, Theorem IX.6.7 in [4]) that if $\varphi \in \mathcal{M}_2(G) = \ell^\infty(\Gamma)$ and $T \in \mathcal{L}(L^2(G))$ satisfy $M_\varphi^{(2)}T = TM_\varphi^{(2)}$, then also $M_{\bar{\varphi}}^{(2)}T = TM_{\bar{\varphi}}^{(2)}$. As noted in the Introduction, this latter result was extended to the case $E = L^p(\mathbb{T})$ with $1 \leq p < \infty$ in Theorem 3.1 of [14]. The purpose of this section to extend this result to all translation and reflection invariant B.f.s.’ with o.c.-norm over arbitrary compact abelian groups (Theorem 5.2).

For any $f \in L^1(G)$ we denote

$$\text{supp}(\hat{f}) = \left\{ \gamma \in \Gamma : \hat{f}(\gamma) \neq 0 \right\}.$$

We begin with the following result.

Lemma 5.1. *Let E be a translation invariant B.f.s. over G with $L^\infty(G) \subseteq E$. Let $T \in \mathcal{L}(E)$ and $\varphi, \psi \in \mathcal{M}_E(G)$.*

- (i) Suppose that $M_\varphi^E T = T M_\psi^E$. Then, for every $\gamma \in \Gamma$, it is the case that $\varphi(\xi) = \psi(\gamma)$ for all $\xi \in \text{supp}((T\gamma)^\wedge)$.
- (ii) Assume, in addition, that E has o.c.-norm. Let φ and ψ have the property that, for each $\gamma \in \Gamma$, we have $\varphi(\xi) = \psi(\gamma)$ for all $\xi \in \text{supp}((T\gamma)^\wedge)$. Then $M_\varphi^E T = T M_\psi^E$.

Proof. (i). Fix $\gamma \in \Gamma$. Since $L^\infty(G) \subseteq E$, we have that $\gamma \in E$ and so

$$M_\varphi^E T\gamma = T M_\psi^E \gamma. \tag{16}$$

Recalling from (12) that $M_\psi^E \gamma = \psi(\gamma)\gamma$, it follows that $T M_\psi^E \gamma = \psi(\gamma) T\gamma$. Hence,

$$(T M_\psi^E \gamma)^\wedge = \psi(\gamma) (T\gamma)^\wedge. \tag{17}$$

On the other hand,

$$(M_\varphi^E T\gamma)^\wedge = \varphi \cdot (T\gamma)^\wedge. \tag{18}$$

A combination of (16), (17) and (18) yields that

$$\varphi(\xi) (T\gamma)^\wedge(\xi) = \psi(\gamma) (T\gamma)^\wedge(\xi), \tag{19}$$

for all $\xi \in \Gamma$. If $\text{supp}((T\gamma)^\wedge) = \emptyset$, then there is nothing to be proved. So, suppose there exists $\xi \in \text{supp}((T\gamma)^\wedge)$. Then $(T\gamma)^\wedge(\xi) \neq 0$ and hence, (19) implies that $\varphi(\xi) = \psi(\gamma)$. This proves part (i).

(ii). Fix $\gamma \in \Gamma$. If $\xi \in \Gamma \setminus \text{supp}((T\gamma)^\wedge)$, i.e., $(T\gamma)^\wedge(\xi) = 0$, then it is clear that (19) holds. On the other hand, if $\xi \in \text{supp}((T\gamma)^\wedge)$, then it follows from the hypothesis that (19) holds. Hence,

$$\varphi \cdot (T\gamma)^\wedge = \psi(\gamma) (T\gamma)^\wedge.$$

Via (17) and (18) this yields that

$$(M_\varphi^E T\gamma)^\wedge = (T M_\psi^E \gamma)^\wedge.$$

The uniqueness of Fourier transforms in $L^1(G)$ then implies (16).

Since $\gamma \in \Gamma$ is arbitrary, the linearity of T , M_φ^E and M_ψ^E imply that

$$M_\varphi^E Tg = T M_\psi^E g, \quad g \in \tau(G),$$

where, as before, $\tau(G) \subseteq L^\infty(G) \subseteq E$ denotes the space of all trigonometric polynomials on G . But, E has o.c.-norm and so $\tau(G)$ is dense in E ; Proposition 4.1. The operators T , M_φ^E and M_ψ^E are continuous and hence, we can conclude that $M_\varphi^E T = T M_\psi^E$. The proof is thereby complete. ■

The following Fuglede type theorem is a consequence of Lemma 5.1.

Theorem 5.2. *Let $E \neq \{0\}$ be a translation and reflection invariant B.f.s. over G with o.c.-norm. Suppose that $\varphi, \psi \in \mathcal{M}_E(G)$ and $T \in \mathcal{L}(E)$ satisfy $M_\varphi^E T = T M_\psi^E$. Then $M_\varphi^E T = T M_\psi^E$.*

Proof. The reflection invariance of E ensures that $\bar{\varphi}, \bar{\psi} \in \mathcal{M}_E(G)$ and the order continuity of the norm in E implies that $L^\infty(G) \subseteq E$. By Lemma 5.1(i), the condition $M_\varphi^E T = T M_\psi^E$ implies, for every $\gamma \in \Gamma$, that

$$\varphi(\xi) = \psi(\gamma), \quad \xi \in \text{supp}((T\gamma)^\wedge). \tag{20}$$

It is clear that the functions $\bar{\varphi}$ and $\bar{\psi}$ (in place of φ and ψ , respectively) also satisfy (20). Therefore, via Lemma 5.1 (ii) applied to $\bar{\varphi}, \bar{\psi} \in \mathcal{M}_E(G)$, we can conclude that $M_{\bar{\varphi}}^E T = T M_{\bar{\psi}}^E$. This completes the proof. ■

The following result is a special case of Theorem 5.2.

Corollary 5.3. *Let $E \neq \{0\}$ be any rearrangement invariant B.f.s. over G with o.c.-norm. Suppose that $\varphi, \psi \in \mathcal{M}_E(G)$ and $T \in \mathcal{L}(E)$ satisfy $M_{\varphi}^E T = T M_{\psi}^E$. Then $M_{\varphi}^E T = T M_{\psi}^E$.*

References

- [1] C.D. Aliprantis, O. Burkinshaw, Positive Operators, Academic Press, Orlando-San Diego-New York, 1985.
- [2] C. Bennett, R. Sharpley, Interpolation of Operators, Academic Press, London-New York, 1988.
- [3] S.K. Berberian, Note on a theorem of Fuglede and Putnam, Proc. Amer. Math. Soc. 10 (1959) 175–182.
- [4] J.B. Conway, A Course in Functional Analysis, Graduate Texts in Mathematics, Vol. 96, Springer-Verlag, New York, 1985.
- [5] B. de Pagter, W.J. Ricker, Fourier multipliers and spectral measures in Banach function spaces, Positivity 13 (2009) 225–241.
- [6] J. Diestel, J.J. Jr. Uhl, Vector Measures, Amer. Math. Soc., Providence, 1977.
- [7] N. Dunford, J.T. Schwartz, Linear Operators Part II, Wiley Interscience, New York, 1963.
- [8] B. Fuglede, A commutativity theorem for normal operators, Proc. Natl. Acad. Sci. 36 (1950) 35–40.
- [9] P.R. Halmos, Commutativity and spectral properties of normal operators, Acta Sci. Math. Szeged 12 Pars B (1950) 153–156.
- [10] P.R. Halmos, Measure Theory, Graduate Texts in Mathematics, Vol. 18, Springer Verlag, New York-Heidelberg-Berlin, 1974.
- [11] R. Larsen, An Introduction to the Theory of Multipliers, Springer, Berlin-Heidelberg, 1971.
- [12] W.A.J. Luxemburg, A.C. Zaanen, Notes on Banach function spaces I–XIII, Netherl. Acad. Wetensch. Proc. A66-A67, 1963–1966.
- [13] P. Meyer-Nieberg, Banach Lattices, Springer-Verlag, Berlin-Heidelberg-New York, 1991.
- [14] G. Mockenhaupt, W.J. Ricker, Fuglede’s theorem, the bicommutant theorem and p -multiplier operators for the circle, J. Oper. Theory 49 (2003) 295–310.
- [15] B. de Pagter, W.J. Ricker, Translation invariant Banach function spaces, vector measures and Fourier multipliers, in preparation.
- [16] C.R. Putnam, On normal operators in Hilbert space, Amer. J. Math. 73 (1951) 357–362.
- [17] H. Radjavi, P. Rosenthal, Invariant Subspaces, Springer-Verlag, New York, 1973.
- [18] M. Rosenblum, On a theorem of Fuglede and Putnam, J. Lond. Math. Soc. 33 (1958) 376–377.
- [19] W. Rudin, Fourier Analysis on Groups, Wiley-Interscience, New York, 1967.
- [20] W. Rudin, Invariant means on L^∞ , Studia Math. 44 (1972) 219–227.
- [21] H.H. Schaefer, Banach Lattices and Positive Operators, Springer-Verlag, Berlin-Heidelberg, 1974.
- [22] A.C. Zaanen, Integration, second ed., North-Holland, Amsterdam, 1967.
- [23] A.C. Zaanen, Riesz Spaces II, North-Holland, Amsterdam, 1983.