

RESEARCH

Open Access



# Professional quality of life of child welfare workers and psychotherapists working with traumatized young unaccompanied refugees in Germany: a cross-sectional study

Pia Maria Schwegler<sup>1\*</sup>, Barbara Kasparik<sup>1</sup>, Jonathan Thielemann<sup>1</sup>, Rebekka Eilers<sup>1</sup>, Elisa Pfeiffer<sup>2,3</sup>, Cedric Sachser<sup>3,4</sup> and Rita Rosner<sup>1</sup>

## Abstract

**Background** Burnout symptoms and secondary traumatic stress are especially high among Child Welfare Workers (CWWs) and psychotherapists and might have increased since the recent increase in refugee numbers. Little is known about the wellbeing of CWW and psychotherapists working with unaccompanied young refugees (UYR), especially in German child and youth welfare facilities where they work closely together. This study aims to assess levels of compassion satisfaction (CS), burnout (BO), and secondary traumatic stress (STS) in German CWWs and psychotherapists, examining connections to demographic variables and group differences.

**Methods**  $N=198$  CWW and  $N=97$  psychotherapists were assessed via the Professional Quality of Life Scale (ProQOL). Descriptive statistics, t-test for independent samples, correlations between the ProQOL scales and hierarchical regression analyses with demographic variables as predictors were computed.

**Results** Both samples showed average to high levels of CS, and low to average levels of BO and STS. CWWs scored significantly higher than psychotherapists on BO and STS. In CWW, demographic variables were not associated with CS, BO or STS. Among psychotherapists, previous experiences in working with UYR ( $\beta=0.38$ ;  $p<.001$ ) were positively associated with CS. Factors such as weekly therapy sessions ( $\beta=-0.35$ ,  $p=.001$ ) and the psychotherapist's migration background ( $\beta=-0.20$ ,  $p=.048$ ) were negatively associated with STS. The number of PTSD cases treated ( $\beta=0.27$ ,  $p=.018$ ) and additional training ( $\beta=0.31$ ,  $p=.006$ ) were positively associated with STS.

**Conclusion** Among psychotherapists, prior experience in working with UYRs may contribute to elevated levels of CS, whereas a higher number of previously treated PTSD cases appears to be associated with increased STS. Further research on the influence of sociodemographic variables is needed for CWWs to identify protective and risk factors. Supporting and training CWWs and psychotherapists is crucial for quality treatment of traumatized UYRs.

**Keywords** Child welfare workers, Psychotherapists, Unaccompanied young refugees, Compassion satisfaction, Burnout, Secondary traumatic stress, ProQOL

\*Correspondence:  
Pia Maria Schwegler  
Pia.schwegler@ku.de

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Introduction

According to the United Nations High Commissioner for Refugees (UNHCR) the number of refugees worldwide has increased over the years (currently 36.4 million) [1]. This includes unaccompanied refugee minors, who are often exposed to traumatic events during their flight [2]. In Germany, child welfare workers (CWW) and psychotherapists work closely together to support UYR.

In general, CWW and psychotherapists belong to occupational groups described as likely to experience psychological distress and emotional burden [3–6]. These stressful experiences may result in feelings of overwhelming responsibility and strained personal relationships [7]. In addition, the workload and high levels of organizational demands have further increased in the last years, and cases are often characterized by complexity and severity, resulting in a high emotional burden for CWW and psychotherapists [8]. Persistent psychological distress can lead to mental and physical exhaustion [9], exit-seeking behaviors and work withdrawal [10]. Moreover, the CWW's daily contact with physically, sexually, and emotionally abused children puts them especially at risk to develop burnout (BO) symptoms, secondary traumatic stress (STS) or a lack of compassion satisfaction (CS) [8, 11–13].

For both, CWW and psychotherapists, a particular challenging field is the work with unaccompanied young refugees (UYR). In Germany, the number of UYR has increased [14] and studies have indicated a higher prevalence of mental disorders among UYR [15–19]. Thus, there is a great need for mental health interventions tailored to the needs of UYR [20]. In the work with refugees, the likelihood of secondary exposure to traumatic events is high for therapists and social workers. Research has shown that BO and STS symptoms are possible consequences of working with a highly traumatized population [21–24]. In their review, Roberts and colleagues [23] reported that individuals working professionally and voluntarily with forcibly displaced people had pooled prevalence rates of 29.7% for high-level BO and 45.7% for moderate to severe-level STS, respectively. In the work with refugees, specific demands include a heightened emotional impact of work, cultural or language barriers and additional administrative obstacles regarding their residence status and other refugee-specific topics [25]. Lastly, working with refugee children seems to be associated with higher STS scores compared to working with refugee adults [21].

For STS, studies with social workers have reported various risk factors, such as female gender, younger age, or years of work experience [8, 12, 26]. Moreover, with increasing years of experience, BO and STS increase while CS seems to decrease. However, an earlier study by Avieli and colleagues [27] showed in a sample of social

workers and therapists that with the increase in years of experience, STS and BO levels decreased significantly, whereas CS levels increased. Thus, in terms of experience, it is unclear whether it should be considered a risk or protective factor and how this may differ depending on the professional role.

Regarding therapists, various risk factors are discussed [28, 29]. Yang and Hayes [30] reported risk factors for BO such as a lack of perceived job control, a demanding caseload, and psychotherapists' mental health such as trauma history. It has also been found that personal trauma experience is a risk factor for compassion fatigue [31]. Moreover, a personal history of trauma and flight experience favors higher levels of STS [21]. Furthermore, Simionato and Simpson [29] found a significant correlation between younger age, having less work experience, and being overinvolved in client problems with moderate to high levels of BO. Regarding STS, some studies have shown that a greater number of patients with PTSD is associated with increased STS [32, 33], while CS appears to increase with work experience [27, 34].

Among professionals who provide therapeutic support to refugees, rapport, and supervision and particularly the alliance with the supervisor was related to well-being, such as professional growth, balance, and employing boundaries [35]. Furthermore, Denkinger and colleagues [21] found that a secure attachment style can function as a protective factor preventing STS in a sample of psychologists, psychotherapists, interpreters, social workers, volunteers, and others who provided treatment and support to refugees from Iraq. Among therapists who provide therapeutic services for traumatized refugees, higher rates of self-efficacy and trauma-specific training led to lower levels of STS [36]. In a study by Plakas and colleagues [37] prior educational level could not be identified as an influencing factor on CS, BO or STS among social workers, coordinators, fieldworkers, or others working with refugees. It has not been investigated whether the therapeutic background (i.e. CBT, depth psychology etc.) has an influence on STS, BO, and CS scores.

Although there appears to be an increase in research that captures the impact of working with refugees in social workers and mental health professionals, there is a lack of knowledge about the wellbeing of CWW and licensed psychotherapists working with UYR. However, this is particularly important as UYR rely more heavily on social workers than accompanied young refugees or refugee adults. Social workers and therapists frequently adopt a more parental position. The aim of the present study is to identify levels of CS, BO, and STS in German CWW and licensed psychotherapists, examine differences between the two groups of professionals, as well as to investigate the influence of demographic and work-related variables on these dimensions. Demographic and

work-related variables vary depending on the respective group. For more detail see Sect. [Statistical analysis](#).

## Materials and methods

### Ethical considerations

Written informed consent was obtained from all individual participants included in the study. The studies involving human participants were reviewed and approved by ethics committees at The Ulm University (No. 243/19) and at The Catholic University of Eichstätt-Ingolstadt (No. 004–19). All data collection is carried out in accordance with the relevant guidelines and regulations following the Declaration of Helsinki.

### Study design and data collection

Participant data were collected between 2020 and 2022 within a greater trial [38]. The main study aims at comparing a stepped care approach to a treatment as usual for UYR. Therefore, psychotherapists were recruited to provide trauma-focused cognitive behavioral therapy [39] to the UYR. The social workers were recruited for external assessments and the implementation of a trauma-focused group intervention “Mein Weg” [40] in the Child and Youth Welfare Service (CYWS) facility. CWWs were recruited through the CYWS facilities where they were employed. Recruitment strategies for CYWS facilities included letters of invitation and telephone contact, digital information events and flyers. A website was created, and press interviews were conducted [38]. Psychotherapists were recruited via email through professional associations and personalized letters. The inclusion criteria for psychotherapist participants were: (1) licensed as either child and adolescent psychotherapists, or psychological psychotherapists, (2) agreement to treat up to three refugees with TF-CBT and (3) comply with all other study regulations. Additional information on recruitment strategies and inclusion criteria can be found in the study protocol [38]. Levels of CS, BO, and STS of CWW and psychotherapist participating in the trial were assessed at baseline. The data presented here corresponds to the baseline assessment before any intervention for the participating UYR had started. Thus, this cross-sectional survey was conducted at a time where some of the psychotherapists may have never worked with refugees but had declared their willingness to treat refugees within the trial. Demographic data were collected using two different standardized questionnaires. There were questions that were only suitable for one group or the other (e.g. for psychotherapists, “On average, how many therapy sessions do you have per week?”).

All participants completed the Professional Quality of Life Scale (ProQOL) [41]. The ProQOL measures the positive and negative effects of working with people who have experienced extremely stressful events. The

self-report inventory contains 30 items on a five-point Likert-Scale (1 = never, 2 = rare, 3 = sometimes, 4 = often, 5 = very often). It consists of three scales: Compassion Satisfaction (10 items, e.g., “I get satisfaction from being able to help people”), Burnout (10 items, e.g., “I feel trapped by my job as a helper”) and Secondary traumatic stress (10 items, e.g., “I feel depressed because of the traumatic experiences of the people I help”). In the present sample, chronbach’s alpha for the scales are satisfactory (CS:  $\alpha=0.88$ , BO:  $\alpha=0.75$ , STS:  $\alpha=0.81$ ). The scales were calculated according to the manual with a higher value on a scale meaning a higher expression of the respective dimension (range 10–50). Recommended cut-off scores are low (0–22), moderate (23–41) and high (42–50) [41].

### Participants

The sample ( $N=374$ ) was composed of two different groups: (1) CWW ( $n=266$ ) who worked in 47 different CYWS facilities, and (2) licensed psychotherapists ( $n=108$ ) participating in the abovementioned trial and working together in the care for UYR. 43 cases of the CWW were excluded from analysis due to a high amount of missing data and 24 cases were excluded due to outliers. Regarding the psychotherapists, 3 cases were excluded due to a high amount of missing data and 8 cases were excluded due to outliers. Missing items refer to the fact that less than 50% or none of the ProQOL items were completed. Moreover, some of the socio-demographic data were completely missing. This left an overall sample of  $n=295$ , CWW ( $n=198$ , age  $M=35.42$ ,  $SD=10.82$ , 72.7% female) and psychotherapists ( $n=97$ , age  $M=44.90$ ,  $SD=8.62$ , 81.4% female). For further information about the participants, see Tables 1 and 2.

### Statistical analyses

All analyses were performed using IBM SPSS Statistics (Version 28.0.0.0). Alpha level of statistical significance was set at  $p<.05$  for all analyses. Outliers were identified using studentized excluded residuals, Cook’s distance, and leverage values [42, 43]. Means and standard deviations were calculated for all variables. Level of CS, BO and STS were also calculated and interpreted using cut-off scores [41]. Partial correlations between ProQOL scales were calculated. Due to the missing prerequisite of normal distribution in the STS score, the correlation with the other scales was performed using Kendall-Tau-b. Bias-corrected and accelerated bootstrap 95% CIs were used. A t-test for independent samples was performed to detect significant differences between the two groups (CWW and psychotherapists) regarding CS, BO, and STS. Additionally, the Mann-Whitney-U test was performed due to the lack of a normal distribution of the STS data from the CWW.

**Table 1** Sociodemographic characteristics of the CWW sample

Characteristics	CWW (N=198)
Age	$M=35.42 (SD=10.82)$
Sex	144 female (72.7%)
Graduation (school)/educational level	95 (48%) GCE 72 (36.4%) A-levels 29 (14.6%) advanced technical college certificate 2 (1%) GCSE O-levels second- ary school leaving certificate
Study or training	111 study (56.1%)
Additional training	40.4%
Work experience in general (years)	$M=8.95 (SD=7.77)$
Work experience in current facility (years)	$M=4.01 (SD=3.54)$
Work experience with UYR (months)	$M=41.87 (SD=32.03)$

Note. GCE A-levels = General Certificate of Education Advanced Level, GCSE O-levels = General Certificate of Secondary Education Ordinary Level, UYR = unaccompanied young refugees

Hierarchical regression analyses were conducted for each of the three factors CS, BO and STS and groups (CWW and psychotherapists). Regression models were estimated for both groups based on previous research findings. The prerequisites for the hierarchical regression analyses were checked and implemented in advance. As

**Table 3** Block wise added variables of the hierarchical regression

Group	Model	Variables included
CWW	Model 1	Age
	Model 2	Gender
	Model 3	Years of work experience, experience with UYRs
	Model 4	Education level, training or academic study, further education
Psychotherapist	Model 1	Age, gender, migration background
	Model 2	Years since license acquisition, previous work with refugees, number of PTSD cases, sessions per week
	Model 3	Training in trauma therapy, schema therapy, hypnotherapy, systemic therapy
	Model 4	Professional background in psychology, social work, teaching, pedagogy, social pedagogy
	Model 5	Therapeutic approaches in depth psychology, behavioral therapy
	Model 6	Work with children and youth, work with adults

presented in Table 3, sociodemographic and work-related variables were included block wise as independent variables in the hierarchical regression analyses.

## Results

Descriptive data of the ProQOL scores as well as the correlations of the ProQOL scales for both groups can be found in Tables 4 and 5.

**Table 2** Sociodemographic characteristics of the psychotherapist sample

Characteristics	Psychotherapists (N=97)
Age	$M=44.90 (SD=8.62)$
Sex	79 female (81.4%)
Migration background	15.5% 3.1% born in a different country 10.3% parents born in a different country 2.1% one parent has another citizenship)
Professional Background	37.1% pedagogy 28.9% psychology 21.6% social work 21.6% social pedagogy 9.3% teaching 92.8% licensed psychotherapist for children and youth 5.2% licensed psychotherapist for adults 5.2% licensed psychotherapist for adults and children/youth Duration in years since the acquisition of the license to practice medicine: $M=7.87 (SD=6.29)$
Treatment clientele/acquisition of the license to practice medicine	
Additional training	70.1% 16.5% systemic therapy, 33% trauma therapy, 10.3% schema therapy, 32% EMDR, 7.2% hypnotherapy
Therapy sessions per week	$M=22.77 (SD=7.72)$
Work experience with UYR	63.9%
Number of PTSD patients that have already been treated	$M=13.68 (SD=15.81)$
Therapeutic approach	Behavioral Therapy=85.6%, Depth Psychology=13.4%, others=1%
Patients to treat	Children and Youth=92.8%, Adults=5.2%, both=5.2%

**Table 4** Descriptive data of the ProQOL scores for CWW and psychotherapists

CWW (n = 198)			Psychotherapists (n = 97)		
	M	SD	Range	Low	Moderate
				High	High
CS	39.72	4.22	29-49	0%	64.6%
BO	21.94	4.18	13-34	59.1%	40.9%
STS	21.17	4.12	13-35	67.2%	32.8%

Note: CS = compassion satisfaction, BO = burnout, STS = secondary traumatic stress

### Differences between CWW and psychotherapists

Regarding the BO scale, there was a significant difference between the BO scores of the psychotherapists ( $M = 19.10$ ;  $SD = 3.55$ ;  $n = 97$ ) as compared to CWW ( $M = 21.94$ ;  $SD = 4.18$ ;  $n = 198$ ), with mean difference of  $-2.84$  (95%-CI  $[-3.81, -1.87]$ ) lower for the psychotherapists,  $t(293) = -5.75$ ,  $p < .001$ . It did represent a medium-sized effect,  $d = -0.71$ . In addition, there was a significant difference between STS of psychotherapists ( $M = 18.12$ ;  $SD = 3.43$ ;  $n = 97$ ) and the CWW ( $M = 21.17$ ;  $SD = 4.12$ ;  $n = 198$ ), with mean difference of  $-3.05$  (95%-CI  $[-4.00, -2.10]$ ) lower for the psychotherapists,  $t(293) = -6.30$ ,  $p < .001$ . It did represent a medium-sized effect,  $d = -0.78$ . The Mann-Whitney-U-test showed that the STS levels in psychotherapists ( $Mdn = 19.10$ ) differ significantly from CWW, ( $Mdn = 21.17$ ),  $U = 5712.00$ ,  $Z = -5.67$ ,  $p < .001$ . There was no significant difference between CS scores of the psychotherapists ( $M = 40.06$ ;  $SD = 4.10$ ;  $n = 97$ ) and the CWW ( $M = 39.72$ ;  $SD = 4.22$ ;  $n = 198$ ), with mean difference of  $0.34$  (95%-CI  $[-0.68, 1.35]$ ),  $t(293) = 0.656$ ,  $p = .513$ ,  $d = 0.08$ .

Additional analysis revealed a significant difference between CS scores of psychotherapists who have worked with refugees before ( $M = 41.24$ ;  $SD = 3.87$ ;  $n = 68$ ) and those who have not ( $M = 37.92$ ;  $SD = 4.21$ ;  $n = 37$ ), with mean difference of  $-3.32$  (95%-CI  $[-4.93, -1.71]$ ),  $t(103) = -4.07$ ,  $p < .001$ ,  $d = -0.83$ . There were no significant differences regarding BO and STS. Comparing CS scores of psychotherapists with therapy experience working with refugees and CWW, CWW reported significant lower CS levels with mean difference of  $-1.42$  (95%-CI  $[-2.62, -0.23]$ ),  $t(258) = 0.654$ ,  $p = .02$ ,  $d = -0.34$ .

### Results of hierarchical regression analysis regarding demographic variables and ProQOL subscales

Hierarchical regression analyses for CWW can be found in Table S1 (additional file 1). All predictors were non-significant.

Detailed information on the models of the hierarchical regression for psychotherapists can be found in the additional file 1 (Table S2). Regarding the association between demographic variables and ProQOL subscales among psychotherapists, model 3 was able to explain the greatest significant amount (27%) of variance in CS [ $F(13, 83) = 2.35$ ,  $p = .01$ ]. The model revealed previous work with UYR as a significant predictor for CS ( $\beta = 0.34$ ;  $p = .002$ ).

Furthermore, model 6 was able to explain the greatest significant amount (39.4%) of variance in STS [ $F(22, 74) = 2.18$ ,  $p = .007$ ]. The analysis revealed that a higher number of weekly therapy sessions was associated with lower levels of STS ( $\beta = -0.35$ ,  $p = .002$ ). Treating more cases diagnosed with PTSD was associated with higher levels of STS ( $\beta = 0.26$ ,  $p = .041$ ). Participants who had

**Table 5** Correlations (r) of the ProQOL scores for CWW and psychotherapists

CWW (n = 198)			Psychotherapists (n = 97)		
	CS	BO	STS	CS	BO
CS	1	−0.475** [−0.571, −0.374]	−0.110* [−0.220, −0.010]	1	−0.524** [−0.660, −0.376]
BO	−0.475**	1	0.363** [0.268, 0.448]	−0.524**	1
STS	−0.110* <sup>a</sup>	0.363*** <sup>a</sup>	1	−0.200*	0.514**

Note. \*\*p < .001 und \*p < .05. BCa bootstrap 95% CIs reported in brackets. CS = compassion satisfaction, BO = burnout, STS = secondary traumatic stress. <sup>a</sup> correlation was performed using Kendall-Tau-b

undergone additional training with a focus on trauma therapy reported higher STS levels ( $\beta = 0.34$ ,  $p = .009$ ). Participants with a migration background reported less STS ( $\beta = -0.20$ ,  $p = .05$ ). There was no significant predictor for BO (see additional file 1, Table S2).

## Discussion

The aim of the present study was to identify levels of CS, BO, and STS in German CWW who work with UYR and licensed psychotherapists and to determine whether there was a difference between these two groups. Furthermore, we investigated whether demographic variables were associated with different levels in the ProQOL dimensions. In our overall sample, the participants reported average levels of BO and STS. Previous studies reported more average- to high-level scores on BO and STS among caregivers, refugee relief workers, professionals, and volunteers [21–24], whereas in our samples, mainly below average to average and no above average scores on BO and STS were found. The CS scores ranged from average to above average. Given that participants were likely already highly interested in working with refugees, CS levels may be elevated. Consequently, the sample mean may exceed that of the general population and should be taken into account when interpreting the results. Furthermore, differences in BO and STS scores between this and other studies may be due to differences in the areas of work. For example, working in receptive and transit centers [24] or working exclusively with women and children who may have experienced sexualized violence [21]. The social workers in our study worked in residential youth groups and the psychotherapists treated UYR alongside people without experience of flight and trauma. Additionally, 36.1% of therapists had not worked with refugees by the time the data was collected. Based on this study, future research should distinguish between different professional groups and areas of work with refugees.

Although the difference was small, psychotherapists scored significantly lower on the BO and STS scales than CWW. Little is known about the differences between professional groups working with refugees or UYR. Živanović and Marković [24] could not find any difference between occupational groups providing different

types of services to refugees, i.e. legal, psychological, medical, or translational services. However, it seems that occupational stress is higher in CWW than in other professions due to various factors like higher caseloads, time pressure, being short staffed, working overtime, administrative regulations and protecting child safety [3, 4]. In our sample, the reason for the differences could be the nature and intensity of the contact. While CWW accompany children and young people in their everyday lives for several hours a day, up to 5 days a week, psychotherapists only see their clients in one to two hour sessions on a weekly basis. Another possibility are differences in training. Therapists learn about the treatment of PTSD during their undergraduate and postgraduate training, whereas social workers receive less intensive training. However, being trained in therapies with a trauma-focus was associated with higher STS in our sample. It should also be noted that our group of psychotherapists had not all worked with UYR or carried out trauma therapy at the time of the analysis. Conversely, comparing therapists who have worked with refugees to those who have not revealed no differences in BO or STS scores. The analysis revealed that therapists with experience treating refugees had higher CS scores than those without. This suggests a positive effect associated with this work. These therapists also scored significantly higher on CS than the CWW group, which may reflect professional differences or statistical effects due to the groups' different sample sizes. If these are professional differences, adapting labor policies to address challenges such as overtime, staff shortages, and high case numbers [3, 4] may reduce BO and STS while improving CS among CWW. However, as this is a cross-sectional survey, the generalizability of the results is limited. A longitudinal study is necessary to observe changes in CS, BO, and STS over time.

Surprisingly, there were no significant predictors in the hierarchical regression for CWW in all models. This finding is contrary to earlier studies that found that female gender and younger age are risk factors for BO and STS [8, 12]. However, in terms of work experience, our results fit the literature as some studies suggest that it puts CWW at risk [8, 26], while others indicate it as a protective factor [27]. When considering a floor effect, given the somewhat lower scores on the ProQOL scales in our

sample, it may be that the gender and age did not come into effect. It may also be necessary to consider variables other than those we included. One possible predictor could be personal trauma history, as other studies have shown that CWW's own trauma history has an impact on CS and STS [21, 31]. Furthermore, we included the predictors work experience in years and in the current facility. However, it might be more important to consider working hours, shift times or overtime, as these may better represent the workload that could be experienced as a burden or cause emotional distress [6]. It is possible that CWW seek intervisionary support from their colleagues. Support from team members has been reported to be essential for well-being [35] and may be a protective factor not captured by the variables in our study.

Among psychotherapists, hierarchical regression results indicated that higher numbers of weekly therapy sessions were associated with lower levels of STS. In this regard, studies are inconsistent, as in some, a positive correlation between the proportion of time spent with traumatized clients and the development of STS was shown [21], while in others, the opposite [44] or no correlation [24] was found. It should be noted that in the present study, we only asked about the number of therapy sessions per week, not specific to PTSD.

The observed association between the number of PTSD cases treated and increased STS scores warrants further investigation. It seems that it is not necessarily the number of patients per week that increases the STS score, but rather the total number of PTSD cases that are treated. These findings are consistent with those of Hensel and colleagues [33], who indicated that the proportion of time spent working with traumatized patients may be more important than the actual number of patients. When selecting cases, psychotherapists should consider this connection and arrange for parallel supervision when treating a high volume of trauma patients. Furthermore, future research should more rigorously collect data on the proportion of traumatized clients treated, trauma characteristics and treatment approaches to further evaluate this aspect.

Unexpectedly, participants who had undergone additional training focused on trauma therapy reported higher levels of STS. Earlier research indicated that the specific trauma training or more education should lead to a lower burden [44]. It is possible that psychotherapists with trauma training also have greater interest in treating PTSD and therefore take on more PTSD cases on average. This explanation aligns with our result that a higher PTSD caseload was associated with more STS. However, the possibility of habituation is important to consider. Additionally, it is unclear how much supervision psychotherapist with advanced trauma training still received which may have an impact on their STS scores [35]. The

literature agrees on the need for trauma training, education, and supervision in the field of working with refugees [21, 24, 31, 35]. Moreover, continued supervision remains essential even for experienced psychotherapists who have treated a higher number of trauma cases. It should not be assumed that increased clinical experience reduces the need for supervision.

Another result was that psychotherapists with a migrant background reported lower STS scores. One potential explanation is that language barriers or cultural differences are less pronounced, leading to greater understanding and openness. In addition, their own migration history could also lead to higher resilience and thus lower STS values. However, Denkinger and colleagues found that a person's own history of flight is a risk factor for STS [21]. Therefore, flight experience and migration background must be considered separately. Furthermore, psychotherapists with a migration or refugee background should receive more support. Although their experiences could offer valuable insights to improve care for UYR, they remain underrepresented in research and should be more actively included.

Furthermore, the analysis showed that greater experience in working with refugees was associated with higher levels of CS. Additionally, there are no significant differences in BO and STS based on whether or not psychotherapists have previously treated refugees. This finding is not consistent with the findings of Mavratza and colleagues [31] who found that working with migrants had a negative impact on professional quality of life, which they attributed to a possible influence on the BO and compassion fatigue scales. Plakas [37] was also able to show that an increase in work experience in humanitarian aid was associated with an increase in BO and a decrease in CS. However, it should be noted that these studies did not exclusively survey psychotherapists but also social workers, nurses, nursing assistants, coordinators, field workers and more. Furthermore, it is possible that psychotherapists with a higher CS trait are more likely to work with refugees. However, these findings suggest that working with refugees can be professionally rewarding. Early-career and experienced psychotherapists alike should be encouraged to engage in this field and provided with the necessary support.

### Limitations

This study has some limitations. First, the sample size varied substantially depending on the group. Subsequently, some of the analyses contained only a few participants and missed statistical power. In addition, some of the psychotherapists (36.1%) had not yet worked with refugees at the time of the survey. To ensure generalizability, future research should survey only therapists who already have experience in working with UYR. Moreover,

questions should have been asked about personal trauma histories and personal experiences of flight. Additionally, we did not assess supervisory support in CWW or psychotherapists. This may be necessary for obtaining further information on work-related stress factors.

Another aspect affecting the sample is the context of the project in which the data were collected. It aims to implement a low-threshold care service for UYR. The CYWS facilities as well as the psychotherapists who register with the project have a certain interest and motivation to work with this special cohort. Therefore, the possibility of a selection bias cannot be ruled out. Furthermore, the data is cross-sectional, whereas longitudinal data could have provided more accurate information on the development of professional quality of life in the work with UYR. Due to the explorative approach, the results can only be generalized to a limited extent. This should be considered when interpreting our results and reviewing them in future studies.

## Conclusion and implications

In sum, BO and STS were present in CWW and psychotherapists, but levels were lower than expected. Psychotherapists had lower STS and BO scores than CWW, suggesting greater exposure to stress for CWW. However, no significant predictors could be detected in the hierarchical regression of the CWW. Among psychotherapists, a greater number of PTSD cases was associated with higher STS scores. It has been shown that a greater number of weekly therapy sessions was linked to lower STS scores. Psychotherapists with trauma training reported higher levels of STS. Having a migration background was associated with lower STS values. Furthermore, psychotherapists with experience in working with refugees reported higher levels of CS than those without.

Given the aforementioned limitations, several implications for future research and professional development emerge. To ensure the quality of treatment and care, it is important to support and train CWW and psychotherapists working with refugees. By doing so, potential barriers or fears of treatment can be removed. This includes adequate trauma-specific training as well as training on the topic of flight in general. Training should be provided not only in clinical settings but also in pedagogical and educational areas to equip young adults in social professions with basic knowledge. Additionally, studies often reference the added stress of working with refugees. However, it is possible that working with UYR can also be fulfilling and not solely diminish professional quality of life. Future research should address the role of occupational groups, treatment approach, and migration background of the psychotherapist. Furthermore, identifying risk and protective factors among CWW is important. Predictors such as trauma history, workload,

and the importance of supervision or intervision support should be considered. Research in the field of professionals working with UYR should increase as it could support professionals' quality of life and ultimately benefit their patients.

## Abbreviations

BO	Burnout
CS	Compassion Satisfaction
CWW	Child Welfare Workers
CYWS	Child and Youth Welfare Service
proQOL	Professional Quality of Life Scale
PTSD	Posttraumatic Stress Disorder
STS	Secondary Traumatic Stress
UYR	Unaccompanied Young Refugees

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13034-025-00942-0>.

Supplementary Material 1: Table S1 and S2: Coefficient of the hierarchical regression analyses (CWW & psychotherapists).

## Acknowledgements

We thank the CWW and psychotherapists for their participation in the survey and their everyday work helping young people.

## Author contributions

The authors' responsibilities were as follows: The BETTER CARE project was designed and implemented by R.R., E.P. and C.S. The data were collected by P.S., B.K., J.T., E.P. and C.S. P.S. and B.K. performed the statistical analysis. P.S. and J.T. drafted the manuscript. B.K., J.T., R.E., E.P., C.S. and R.R. critically reviewed the manuscript. The final supervision was carried out by R.R. All authors read and approved the final manuscript.

## Funding

Open Access funding enabled and organized by Projekt DEAL. This work is part of the BETTER CARE project supported by the German Ministry of Education and Research (01EF1802A, 01EF1802B).

## Data availability

Our data were collected by the research group and are not publicly available. The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Declarations

### Ethics approval and consent to participate

Written informed consent was obtained from all individual participants included in the study. The studies involving human participants were reviewed and approved by ethics committees at The Ulm University (No. 243/19) and at The Catholic University of Eichstätt-Ingolstadt (No. 004-19). All data collection is carried out in accordance with the relevant guidelines and regulations following the Declaration of Helsinki.

### Consent for publication

Not applicable.

### Competing interests

The authors declare no competing interests.

### Author details

<sup>1</sup>Department of Clinical and Biological Psychology, Catholic University Eichstätt-Ingolstadt, Eichstätt-Ingolstadt 85072, Germany

<sup>2</sup>Department of Clinical Psychology and Child and Adolescent Psychotherapy, Catholic University Eichstätt-Ingolstadt, Eichstätt-Ingolstadt, Germany

<sup>3</sup>Department of Child and Adolescent Psychiatry/Psychotherapy, Ulm University, Ulm 89070, Germany

<sup>4</sup>Department of Clinical Child and Adolescent Psychology, Institute of Psychology, University of Bamberg, Bamberg, Germany

Received: 8 April 2025 / Accepted: 2 July 2025

Published online: 15 July 2025

## References

- United Nations High Commissioner for Refugees. Refugee Data Finder [Internet]. UNHCR. 2023 [updated 2023 Oktober 24; cited 2023 November 26]. Available from: <https://www.unhcr.org/refugee-statistics/>
- World Health Organization. Refugee and migrant health [Internet]. WHO. 2023 [updated 2022; cited 2023 November 26]. Available from: <https://www.who.int/news-room/fact-sheets/detail/refugee-and-migrant-health#:~:text=Key%20facts,4.4%20million%20are%20asylum%20seekers>
- Wilke DJ, Randolph K, Olson C. Examining occupational stress in early-career child welfare workers. *J Workplace Behav Health*. 2020;158–74. <https://doi.org/10.1080/15555240.2020.1807354>.
- Poulsen I. Weichen Neu stellen! Ein Praxishandbuch Zur selbstfürsorge und burnoutprävention für fachkräfte in sozialen und pädagogischen Berufen. 2nd ed. Books on Demand; 2018.
- Nissen-Lie HA, Orlinsky DE, Rønnestad MH. The emotionally burdened psychotherapist: personal and situational risk factors. *Prof Psychology: Res Pract*. 2021;52:429–38. <https://doi.org/10.1037/pro0000387>.
- Hermon SR, Chahla R. A longitudinal study of stress and satisfaction among child welfare workers. *J Social Work*. 2018;192–215. <https://doi.org/10.1177/468017318757557>.
- Råbu M, Moltu C, Binder PE, McLeod J. How does practicing psychotherapy affect the personal life of the therapist? A qualitative inquiry of senior therapists' experiences. *Psychother Res*. 2016;26:737–49. <https://doi.org/10.1080/0503307.2015.1065354>.
- Salloum A, Kondrat DC, Johnco C, Olson KR. The role of self-care on compassion satisfaction, burnout and secondary trauma among child welfare workers. *Child Youth Serv Rev*. 2015;49:54–61. <https://doi.org/10.1016/j.childyouth.2014.12.023>.
- Meyers TW, Cornille TA. The trauma of working with traumatized children. *Treat Compassion Fatigue*, 1st edn. London: Routledge; 2002; pp 39–55.
- Travis DJ, Lizano EL, Mor Barak ME. I'm so stressed!: A longitudinal model of stress, burnout and engagement among social workers in child welfare settings. *Br J Social Work*. 2016;46:1076–95. <https://doi.org/10.1093/bjsw/bct205>.
- Rienks SL. An exploration of child welfare caseworkers' experience of secondary trauma and strategies for coping. *Child Abuse Negl*. 2020;110:104355. <https://doi.org/10.1016/j.chab.2020.104355>.
- Molnar BE, Meeker SA, Manners K, Tieszen L, Kalergis K, Fine JE, et al. Vicarious traumatization among child welfare and child protection professionals: A systematic review. *Child Abuse Negl*. 2020;110:104679. <https://doi.org/10.1016/j.chab.2020.104679>.
- Letson MM, Davins C, Sherfield J, Beer OWJ, Phillips R, Wolf KG. Identifying compassion satisfaction, burnout, & traumatic stress in children's advocacy centers. *Child Abuse Negl*. 2020;110:104240. <https://doi.org/10.1016/j.chab.2019.104240>.
- Bundesamt für Migration und Flüchtlinge. Aktuelle Zahlen [internet]. BMF. 2022 [updated 2022 March 7; cited 2023 June 2]. Available from: [https://www.bamf.de/SharedDocs/Anlagen/DE/Statistik/AsylinZahlen/aktuelle-zahlen-februar-2022.pdf?\\_\\_blob=publicationFile&v=3](https://www.bamf.de/SharedDocs/Anlagen/DE/Statistik/AsylinZahlen/aktuelle-zahlen-februar-2022.pdf?__blob=publicationFile&v=3)
- Acarturk C, McGrath M, Roberts B, Ilkkursun Z, Cuijpers P, Sijbrandij M, et al. Prevalence and predictors of common mental disorders among Syrian refugees in istanbul, turkey: a cross-sectional study. *Soc Psychiatry Psychiatr Epidemiol*. 2021;56:475–84. <https://doi.org/10.1007/s00127-020-01941-6>.
- Biddle L, Menold N, Bentner M, Nöst S, Jahn R, Ziegler S, Bozorgmehr K. Health monitoring among asylum seekers and refugees: a state-wide, cross-sectional, population-based study in Germany. *Emerg Themes Epidemiol*. 2019;16:3. <https://doi.org/10.1186/s12982-019-0085-2>.
- Müller LRF, Büter KP, Rosner R, Unterhitzberger J. Mental health and associated stress factors in accompanied and unaccompanied refugee minors resettled in germany: A cross-sectional study. *Child Adolesc Psychiatry Mental Health*. 2019;13:8. <https://doi.org/10.1186/s13034-019-0268-1>.
- Vervliet M, Lammertyn J, Broekaert E, Derluyn I. Longitudinal follow-up of the mental health of unaccompanied refugee minors. *Eur Child Adolesc Psychiatry*. 2014;23:337–46. <https://doi.org/10.1007/s00787-013-0463-1>.
- Hornfeck F, Eglinsky J, Garbade M, Rosner R, Kindler H, Pfeiffer E, Sachser C. Mental health problems in unaccompanied young refugees and the impact of post-flight factors on PTSS, depression and anxiety—A secondary analysis of the better care study. *Front Psychol*. 2023;14:1149634. <https://doi.org/10.3389/fpsyg.2023.1149634>.
- Baron J, Flory L. Versorgungsbericht zur psychosozialen Versorgung von Flüchtlingen und Folteropfern in Deutschland [internet]. BAFF; 2020 [updated 2020; cited 2023 July 16]. Available from: <https://www.baff-zentren.org/aktuelles/viele-traumatisierte-gefluechte-nicht-versorgt/>
- Denkinger JK, Windhorst P, Rometsch-Ogioun El Sount C, Blume M, Sedik H, Kizilhan JL, et al. Secondary traumatization in caregivers working with women and children who suffered extreme violence by the Islamic state. *Front Psychiatry*. 2018;9:234. <https://doi.org/10.3389/fpsyg.2018.00234>.
- Ebren G, Demircioğlu M, Çırakoğlu OC. A neglected aspect of refugee relief works: secondary and vicarious traumatic stress. *J Trauma Stress*. 2022;35:891–900. <https://doi.org/10.1002/jts.22796>.
- Roberts F, Teague B, Lee J, Rushworth I. The prevalence of burnout and secondary traumatic stress in professionals and volunteers working with forcibly displaced people: A systematic review and two Meta-Analyses. *J Trauma Stress*. 2021;34:773–85. <https://doi.org/10.1002/jts.22659>.
- Živanović M, Marković MV. Latent structure of secondary traumatic stress, its precursors, and effects on people working with refugees. *PLoS ONE*. 2020;15:10e0241545. <https://doi.org/10.1371/journal.pone.0241545>.
- Reißmann S, Flothow A, Harth V, Mache S. Exploring job demands and resources in psychotherapists treating displaced people - A scoping review. *Psychother Res*. 2022;32:1076–89. <https://doi.org/10.1080/10503307.2022.2071653>.
- Weiss-Dagan S, Ben-Porat A, Itzhaky H. Secondary traumatic stress and vicarious post-traumatic growth among social workers who have worked with abused children. *J Social Work*. 2022;22:170–87. <https://doi.org/10.1177/146017320981363>.
- Avieli H, Ben-David S, Levy I. Predicting professional quality of life among professional and volunteer caregivers. *Psychol Trauma: Theory Res Pract Policy*. 2016;8:80–7. <https://doi.org/10.1037/tra0000066>.
- Kotera Y, Maxwell-Jones R, Edwards AM, Knutton N. Burnout in professional psychotherapists: relationships with Self-Compassion, Work-Life balance, and telepresence. *Int J Environ Res Public Health*. 2021;18:5308. <https://doi.org/10.3390/ijerph18105308>.
- Simionato GK, Simpson S. Personal risk factors associated with burnout among psychotherapists: A systematic review of the literature. *J Clin Psychol*. 2018;74:1431–56. <https://doi.org/10.1002/jclp.22615>.
- Yang Y, Hayes JA. Causes and consequences of burnout among mental health professionals: A practice-oriented review of recent empirical literature. *Psychotber (Chic)*. 2020;57:426–36. <https://doi.org/10.1037/pst0000037>.
- Mavratza T, Adamakidou T, Apostolara P, Fasoi G, Mantzorou M, Mantoudi A, et al. Professional quality of life in Greek health professionals working with refugees and migrants. *Mater Sociomed*. 2021;33:94–9. <https://doi.org/10.5455/msm.2021.33.94-99>.
- Canfield J. Secondary traumatization, burnout, and vicarious traumatization. *Smith Coll Stud Social Work*. 2005;75:81–101. [https://doi.org/10.1300/J497v7n02\\_06](https://doi.org/10.1300/J497v7n02_06).
- Hensel JM, Ruiz C, Finney C, Dewa CS. Meta-analysis of risk factors for secondary traumatic stress in therapeutic work with trauma victims. *J Trauma Stress*. 2015;28:83–91. <https://doi.org/10.1002/jts.21998>.
- Craig CD, Sprang G. Compassion satisfaction, compassion fatigue, and burnout in a National sample of trauma treatment therapists. *Anxiety Stress Coping*. 2010;23:319–39. <https://doi.org/10.1080/10615800903085818>.
- Posselt M, Baker A, Deans C, Procter N. Fostering mental health and well-being among workers who support refugees and asylum seekers in the Australian context. *Health Social Care Community*. 2020;28:1658–70. <https://doi.org/10.1111/hsc.12991>.
- Isawi DT, Post PB. Self-Efficacy of counselors working with refugees. *Adulstspan J*. 2020;19:107–17. <https://doi.org/10.1002/adsp.12099>.
- Plakas C, Burnout C, Fatigue, and Secondary Traumatic Stress Among Humanitarian Aid Workers in Jordan [internet]. 2018 [cited 2023 July 16]. Available from: [https://www.researchgate.net/profile/Christina-Plakas-2/publication/328685237\\_Burnout\\_Compassion\\_Fatigue\\_and\\_Secondary\\_Traumatic\\_Stress\\_Among\\_Humanitarian\\_Aid\\_Workers\\_in\\_Jordan/links/62b9d956056](https://www.researchgate.net/profile/Christina-Plakas-2/publication/328685237_Burnout_Compassion_Fatigue_and_Secondary_Traumatic_Stress_Among_Humanitarian_Aid_Workers_in_Jordan/links/62b9d956056)

dae24e8e75757/Burnout-Compassion-Fatigue-and-Secondary-Traumatic-Str  
ess-Among-Humanitarian-Aid-Workers-in-Jordan.pdf

38. Rosner R, Sachser C, Hornfeck F, Kilian R, Kindler H, Muche R, Müller LRF, Thielemann J, Waldmann T, Ziegenhain U, Unterhitzenberger J, Pfeiffer E. Improving mental health care for unaccompanied young refugees through a stepped-care approach versus usual care--: study protocol of a cluster randomized controlled hybrid effectiveness implementation trial. *Trials*. 2020 <https://doi.org/10.1186/s13063-020-04922-x>.
39. Cohen JA, Mannarino AP, Deblinger E. Treating trauma and traumatic grief in children and adolescents. Guilford; 2016.
40. Pfeiffer E, Goldbeck L. Evaluation of a trauma-focused group intervention for unaccompanied young refugees: A pilot study. *J Trauma Stress*. 2017;30:531–6. <https://doi.org/10.1002/jts.22218>.
41. Stamm BH. The Concise ProQOL Manual 2010 [internet]. 2010 [cited 2023 June 2]. Available from: <https://proqol.org/proqol-manual>
42. Salkind N, editor. Encyclopedia of research design. SAGE Publications, Inc; 2010. <https://doi.org/10.4135/9781412961288>.
43. Vellman PF, Welsch RE. Efficient computing of regression diagnostics. *Am Stat*. 1981;234–42. <https://doi.org/10.1080/00031305.1981.10479362>.
44. Baird S, Jenkins SR, Vicarious, Traumatization. Secondary traumatic stress, and burnout in sexual assault and domestic violence agency staff. *Violence Vict*. 2003;18:71–86. <https://doi.org/10.1891/vivi.2003.18.1.71>.

#### Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.